

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

1. NAME OF THE MEDICINAL PRODUCT

CABOMETRYX 20 mg film-coated tablets
CABOMETRYX 40 mg film-coated tablets
CABOMETRYX 60 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

CABOMETRYX 20 mg film-coated tablets

Each film-coated tablet contains cabozantinib (S)-malate equivalent to 20 mg cabozantinib.

Excipients with known effect

Each film-coated tablet contains 15.54 mg lactose.

CABOMETRYX 40 mg film-coated tablets

Each film-coated tablet contains cabozantinib (S)-malate equivalent to 40 mg cabozantinib.

Excipients with known effect

Each film-coated tablet contains 31.07 mg lactose.

CABOMETRYX 60 mg film-coated tablets

Each film-coated tablet contains cabozantinib (S)-malate equivalent to 60 mg cabozantinib.

Excipients with known effect

Each film-coated tablet contains 46.61 mg lactose

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet.

CABOMETRYX 20 mg film-coated tablets

The tablets are yellow round with no score, and debossed with “XL” on one side and “20” on the other side of the tablet.

CABOMETRYX 40 mg film-coated tablets

The tablets are yellow triangle shaped with no score, and debossed with “XL” on one side and “40” on the other side of the tablet.

CABOMETRYX 60 mg film-coated tablets

The tablets are yellow oval shaped with no score, and debossed with “XL” on one side and “60” on the other side of the tablet.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

CABOMETYX is indicated for the treatment of advanced renal cell carcinoma (RCC) in adults following prior vascular endothelial growth factor (VEGF)-targeted therapy

4.2 Posology and method of administration

Therapy with CABOMETYX should be initiated by a physician experienced in the administration of anticancer medicinal products.

Posology

CABOMETYX (cabozantinib) tablets and COMETRIQ (cabozantinib) capsules are not bioequivalent and should not be used interchangeably (see section 5.2). If a patient must switch from cabozantinib capsules to cabozantinib tablets, the patient should continue at a CABOMETYX dose not to exceed 60 mg or the current COMETRIQ dose (whichever is lower).

The recommended dose of CABOMETYX is 60 mg once daily. Treatment should continue until the patient is no longer clinically benefiting from therapy or until unacceptable toxicity occurs.

Management of suspected adverse drug reactions may require temporary interruption and/or dose reduction of CABOMETYX therapy (see Table 1). When dose reduction is necessary, it is recommended to reduce to 40 mg daily, and then to 20 mg daily. Dose interruptions are recommended for management of CTCAE grade 3 or greater toxicities or intolerable grade 2 toxicities. Dose reductions are recommended for events that, if persistent, could become serious or intolerable.

If a patient misses a dose, the missed dose should not be taken if it is less than 12 hours before the next dose.

Table 1: Recommended CABOMETYX dose modifications for adverse reactions

Adverse reaction and severity	Treatment Modification
Grade 1 and Grade 2 adverse reactions which are tolerable and easily managed	Dose adjustment is usually not required. Consider adding supportive care as indicated.
Grade 2 adverse reactions which are intolerable and cannot be managed with a dose reduction or supportive care	Interrupt treatment until the adverse reaction resolves to Grade ≤ 1 . Add supportive care as indicated. Consider re-initiating at a reduced dose.
Grade 3 adverse reactions (except clinically nonrelevant laboratory abnormalities)	Interrupt treatment until the adverse reaction resolves to Grade ≤ 1 . Add supportive care as indicated. Re-initiate at a reduced dose.
Grade 4 adverse reactions (except clinically nonrelevant laboratory abnormalities)	Interrupt treatment. Institute appropriate medical care. If adverse reaction resolves to Grade ≤ 1 , re-initiate at a reduced dose. If adverse reaction does not resolve, permanently discontinue CABOMETYX.

Note: Toxicity grades are in accordance with National Cancer Institute Common Terminology Criteria for Adverse Events Version 4.0 (NCI-CTCAE v4)

Concomitant medicinal products

Concomitant medicinal products that are strong inhibitors of CYP3A4 should be used with caution, and chronic use of concomitant medicinal products that are strong inducers of CYP3A4 should be avoided (see sections 4.4 and 4.5).

Selection of an alternative concomitant medicinal product with no or minimal potential to induce or inhibit CYP3A4 should be considered.

Special populations

Elderly patients

No specific dose adjustment for the use of cabozantinib in older people (≥ 65 years) is recommended.

Race

There is little experience with cabozantinib in non-White patients.

Patients with renal impairment

Cabozantinib should be used with caution in patients with mild or moderate renal impairment. Cabozantinib is not recommended for use in patients with severe renal impairment as safety and efficacy have not been established in this population.

Patients with hepatic impairment

In patients with mild or moderate hepatic impairment the recommended dose is 40 mg once daily. Patients should be monitored for adverse events and dose adjustment or treatment interruption should be considered as needed (see section 4.2). Cabozantinib is not recommended for use in patients with severe hepatic impairment as safety and efficacy have not been established in this population.

Patients with cardiac impairment

There is limited data in patients with cardiac impairment. No specific dosing recommendations can be made.

Paediatric population

The safety and efficacy of cabozantinib in children and adolescents aged <18 years have not yet been established. No data are available.

Method of administration

CABOMETYX is for oral use. The tablets should be swallowed whole and not crushed. Patients should be instructed to not eat anything for at least 2 hours before through 1 hour after taking CABOMETYX.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

As most events can occur early in the course of treatment, the physician should evaluate the patient closely during the first eight weeks of treatment to determine if dose modifications are warranted. Events that generally have early onset include hypocalcaemia, hypokalaemia, thrombocytopenia, hypertension, palmar-plantar erythrodysesthesia syndrome (PPES), proteinuria, and gastrointestinal (GI) events (abdominal pain, mucosal inflammation, constipation, diarrhoea, vomiting).

Dose reductions and dose interruptions due to an AE occurred in 59.8% and 70%, respectively, of cabozantinib-treated patients in the pivotal clinical trial. Two dose reductions were required in 19.3% of patients. The median time to first dose reduction was 55 days, and to first dose interruption was 38 days.

Perforations and fistulas

Serious gastrointestinal (GI) perforations and fistulas, sometimes fatal, have been observed with cabozantinib. Patients who have inflammatory bowel disease (e.g., Crohn's disease, ulcerative colitis, peritonitis, diverticulitis, or appendicitis), have tumour infiltration in the GI tract, or have complications from prior GI surgery (particularly when associated with delayed or incomplete healing) should be carefully evaluated before initiating cabozantinib therapy and subsequently they should be monitored closely for symptoms of perforations and fistulas including abscesses. Persistent or recurring diarrhoea while on treatment may be a risk factor for the development of anal fistula. Cabozantinib should be discontinued in patients who experience a GI perforation or a fistula that cannot be adequately managed .

Thromboembolic events

Events of venous thromboembolism, including pulmonary embolism, and events of arterial thromboembolism have been observed with cabozantinib. Cabozantinib should be used with caution in patients who are at risk for, or who have a history of, these events. Cabozantinib should be discontinued in patients who develop an acute myocardial infarction or any other clinically significant arterial thromboembolic complication.

Haemorrhage

Severe haemorrhage has been observed with cabozantinib. Patients who have a history of severe bleeding prior to treatment initiation should be carefully evaluated before initiating cabozantinib therapy. Cabozantinib should not be administered to patients that have or are at risk for severe haemorrhage.

Wound complications

Wound complications have been observed with cabozantinib. Cabozantinib treatment should be stopped at least 28 days prior to scheduled surgery, including dental surgery, if possible. The decision to resume cabozantinib therapy after surgery should be based on clinical judgment of adequate wound healing. Cabozantinib should be discontinued in patients with wound healing complications requiring medical intervention.

Hypertension

Hypertension has been observed with cabozantinib. Blood pressure should be well-controlled prior to initiating cabozantinib. During treatment with cabozantinib, all patients should be monitored for hypertension and treated as needed with standard anti-hypertensive therapy. In the case of persistent hypertension despite use of anti-hypertensives, the cabozantinib dose should be reduced. Cabozantinib should be discontinued if hypertension is severe and persistent despite anti-hypertensive therapy and dose reduction of cabozantinib. In case of hypertensive crisis, cabozantinib should be discontinued.

Palmar-plantar erythrodysaesthesia syndrome

Palmar-plantar erythrodysaesthesia syndrome (PPES) has been observed with cabozantinib. When PPES is severe, interruption of treatment with cabozantinib should be considered. Cabozantinib should be restarted with a lower dose when PPES has been resolved to grade 1.

Proteinuria

Proteinuria has been observed with cabozantinib. Urine protein should be monitored regularly during cabozantinib treatment. Cabozantinib should be discontinued in patients who develop nephrotic syndrome.

Reversible posterior leukoencephalopathy syndrome

Reversible Posterior Leukoencephalopathy Syndrome (RPLS), also known as Posterior Reversible Encephalopathy Syndrome (PRES), has been observed with cabozantinib. This syndrome should be considered in any patient presenting with multiple symptoms, including seizures, headache, visual disturbances, confusion or altered mental function. Cabozantinib treatment should be discontinued in patients with RPLS.

Prolongation of QT interval

Cabozantinib should be used with caution in patients with a history of QT interval prolongation, patients who are taking antiarrhythmics, or patients with relevant pre-existing cardiac disease, bradycardia, or electrolyte disturbances. When using cabozantinib, periodic monitoring with on-treatment ECGs and electrolytes (serum calcium, potassium, and magnesium) should be considered.

CYP3A4 inducers and inhibitors

Cabozantinib is a CYP3A4 substrate. Concurrent administration of cabozantinib with the strong CYP3A4 inhibitor ketoconazole resulted in an increase in cabozantinib plasma exposure. Caution is required when administering cabozantinib with agents that are strong CYP3A4 inhibitors. Concurrent administration of cabozantinib with the strong CYP3A4 inducer rifampicin resulted in a decrease in cabozantinib plasma exposure. Therefore chronic administration of agents that are strong CYP3A4 inducers with cabozantinib should be avoided (see sections 4.2 and 4.5).

P-glycoprotein substrates

Cabozantinib was an inhibitor ($IC_{50} = 7.0 \mu M$), but not a substrate, of P-glycoprotein (P-gp) transport activities in a bi-directional assay system using MDCK-MDR1 cells. Therefore, cabozantinib may have the potential to increase plasma concentrations of co-administered substrates of P-gp. Subjects should be cautioned regarding taking a P-gp substrate (e.g., fexofenadine, aliskiren, ambrisentan, dabigatran etexilate, digoxin, colchicine, maraviroc, posaconazole, ranolazine, saxagliptin, sitagliptin, talinolol, tolvaptan) while receiving cabozantinib (see section 4.5).

MRP2 inhibitors

Administration of MRP2 inhibitors may result in increases in cabozantinib plasma concentrations. Therefore, concomitant use of MRP2 inhibitors (e.g. cyclosporine, efavirenz, emtricitabine) should be approached with caution (see section 4.5).

Excipient related warnings

Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

Effect of other medicinal products on cabozantinib

CYP3A4 inhibitors and inducers

Administration of the strong CYP3A4 inhibitor ketoconazole (400 mg daily for 27 days) to healthy volunteers decreased cabozantinib clearance (by 29%) and increased single-dose plasma cabozantinib exposure (AUC) by 38%. Therefore co-administration of strong CYP3A4 inhibitors (e.g., ritonavir, itraconazole, erythromycin, clarithromycin, grapefruit juice) with cabozantinib should be approached with caution.

Administration of the strong CYP3A4 inducer rifampicin (600 mg daily for 31 days) to healthy volunteers increased cabozantinib clearance (4.3-fold) and decreased single-dose plasma cabozantinib exposure (AUC) by 77%. Chronic co-administration of strong CYP3A4 inducers (e.g., phenytoin, carbamazepine, rifampicin, phenobarbital or herbal preparations containing St. John's Wort [*Hypericum perforatum*]) with cabozantinib should therefore be avoided.

Gastric pH modifying agents

Co-administration of proton pump inhibitor (PPI) esomeprazole (40 mg daily for 6 days) with a single dose of 100 mg cabozantinib to healthy volunteers resulted in no clinically-significant effect on plasma cabozantinib exposure (AUC). No dose adjustment is indicated when gastric pH modifying agents (i.e., PPIs, H₂ receptor antagonists, and antacids) are co-administered with cabozantinib.

MRP2 inhibitors

In vitro data demonstrate that cabozantinib is a substrate of MRP2. Therefore, administration of MRP2 inhibitors may result in increases in cabozantinib plasma concentrations.

Bile salt-sequestering agents

Bile salt-sequestering agents such as cholestyramine and cholestagel may interact with cabozantinib and may impact absorption (or reabsorption) resulting in potentially decreased exposure (see section 5.2). The clinical significance of these potential interactions is unknown.

Effect of cabozantinib on other medicinal products

The effect of cabozantinib on the pharmacokinetics of contraceptive steroids has not been investigated. As unchanged contraceptive effect may not be guaranteed, an additional contraceptive method, such as a barrier method, is recommended.

Because of high plasma protein binding levels of cabozantinib (section 5.2) a plasma protein displacement interaction with warfarin may be possible. In case of such combination, INR values should be monitored.

P-glycoprotein substrates

Cabozantinib was an inhibitor (IC₅₀ = 7.0 μM), but not a substrate, of P-gp transport activities in a bi-directional assay system using MDCK-MDR1 cells. Therefore, cabozantinib may have the potential to increase plasma concentrations of co-administered substrates of P-gp. Subjects should be cautioned regarding taking a P-gp substrate (e.g., fexofenadine, aliskiren, ambrisentan, dabigatran etexilate, digoxin, colchicine, maraviroc, posaconazole, ranolazine, saxagliptin, sitagliptin, talinolol, tolvaptan) while receiving cabozantinib.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential/Contraception in males and females

Women of childbearing potential must be advised to avoid pregnancy while on cabozantinib. Female partners of male patients taking cabozantinib must also avoid pregnancy. Effective methods of contraception should be used by male and female patients and their partners during therapy, and for at least 4 months after

completing therapy. Because oral contraceptives might possibly not be considered as “effective methods of contraception”, they should be used together with another method, such as a barrier method (see section 4.5).

Pregnancy

There are no studies in pregnant women using cabozantinib. Studies in animals have shown embryo-foetal and teratogenic effects (see section 5.3). The potential risk for humans is unknown. Cabozantinib should not be used during pregnancy unless the clinical condition of the woman requires treatment with cabozantinib.

Breast-feeding

It is not known whether cabozantinib and/or its metabolites are excreted in human milk. Because of the potential harm to the infant, mothers should discontinue breast-feeding during treatment with cabozantinib, and for at least 4 months after completing therapy.

Fertility

There are no data on human fertility. Based on non-clinical safety findings, male and female fertility may be compromised by treatment with cabozantinib (see section 5.3). Both men and women should be advised to seek advice and consider fertility preservation before treatment.

4.7 Effects on ability to drive and use machines

Cabozantinib has minor influence on the ability to drive and use machines. Adverse reactions such as fatigue and weakness have been associated with cabozantinib. Therefore, caution should be recommended when driving or operating machines.

4.8 Undesirable effects

Summary of safety profile

The most common serious adverse reactions associated with cabozantinib are abdominal pain (3%), pleural effusion (3%), diarrhoea (2%), and nausea (2%). The most frequent adverse reactions of any grade (experienced by at least 25% of patients) included diarrhoea (74%), fatigue (56%), nausea (50%), decreased appetite (46%), palmar-plantar erythrodysesthesia syndrome (PPES) (42%), hypertension (37%), vomiting (32%), weight decreased (31%), and constipation (25%).

Tabulated list of adverse reactions

Adverse reactions are listed in Table 2 according to MedDRA system organ class and frequency categories. Frequencies are based on all grades and defined as: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Table 2: Adverse reactions reported with cabozantinib

MedDRA System Organ Class	Very Common	Common	Uncommon
Infections and infestations		abscess	
Blood and lymphatic disorders	anaemia		
Endocrine disorders	hypothyroidism		
Metabolism and nutrition disorders	decreased appetite, hypophosphataemia, hypoalbuminaemia, hypomagnesaemia, hyponatraemia, hypokalaemia, hyperkalaemia, hypocalcaemia, hyperbilirubinemia	dehydration	
Nervous system disorders	dysgeusia, headache, dizziness		convulsion
Ear and labyrinth disorders		tinnitus	
Vascular disorders	hypertension	pulmonary embolism	
Respiratory, thoracic, and mediastinal disorders	dysphonia, dyspnoea, cough		
Gastrointestinal disorders	diarrhoea, nausea, vomiting, stomatitis, constipation, abdominal pain, dyspepsia	abdominal pain upper, gastrooesophageal reflux disease, haemorrhoids	anal fistula, pancreatitis
Hepatobiliary disorders			hepatitis cholestatic
Skin and subcutaneous tissue disorders	palmar-plantar erythrodysesthesia syndrome, rash, dry skin	pruritus, alopecia	
Musculoskeletal and connective tissue disorders	pain in extremity, muscle spasms, arthralgia		osteonecrosis of the jaw
Renal and urinary disorders	proteinuria		
General disorders and administration site conditions	fatigue, mucosal inflammation, asthenia	peripheral oedema	

MedDRA System Organ Class	Very Common	Common	Uncommon
Investigations	weight decreased, serum ALT, AST, and ALP increased, creatinine increased, triglycerides increased, hyperglycaemia, hypoglycaemia, lymphopenia, neutropenia, thrombocytopenia, GGT increased, amylase increased, blood cholesterol increased, lipase increased		

Description of selected adverse reactions

Data for the following reactions are based on patients who received Cabometyx 60 mg qd po in the pivotal RCC study (section 5.1).

Gastrointestinal (GI) perforation

GI perforations were reported in 0.9% of cabozantinib-treated RCC patients (3/331). Events were Grade 2 or 3. Median time to onset was 10.0 weeks. Fatal perforations have occurred in the cabozantinib clinical program.

Fistulas

Fistulas were reported in 1.2% (4/331) of cabozantinib-treated patients, and included anal fistulas in 0.6% (2/331) cabozantinib-treated patients. One event was Grade 3; the remainder was Grade 2. Median time to onset was 30.3 weeks.

Haemorrhage

The incidence of severe haemorrhagic events (Grade \geq 3) was 2.1% in cabozantinib-treated RCC patients (7/331). Median time to onset was 20.9 weeks. Fatal haemorrhages have occurred in the cabozantinib clinical program.

Reversible Posterior Leukoencephalopathy Syndrome (RPLS)

No case of RPLS was reported in this study, but RPLS has been reported in other clinical studies.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via [the national reporting system listed in Appendix V](#).

4.9 Overdose

There is no specific treatment for cabozantinib overdose and possible symptoms of overdose have not been established.

In the event of suspected overdose, cabozantinib should be withheld and supportive care instituted. Metabolic clinical laboratory parameters should be monitored at least weekly or as deemed clinically appropriate to assess any possible changing trends. Adverse reactions associated with overdose are to be treated symptomatically.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: antineoplastic agent, protein kinase inhibitor, ATC code: L01XE26.

Mechanism of action

Cabozantinib is a small molecule that inhibits multiple receptor tyrosine kinases (RTKs) implicated in tumour growth and angiogenesis, pathologic bone remodeling, drug resistance, and metastatic progression of cancer. Cabozantinib was evaluated for its inhibitory activity against a variety of kinases and was identified as an inhibitor of MET (hepatocyte growth factor receptor protein) and VEGF (vascular endothelial growth factor) receptors. In addition, cabozantinib inhibits other tyrosine kinases including the GAS6 receptor (AXL), RET, ROS1, TYRO3, MER, the stem cell factor receptor (KIT), TRKB, Fms-like tyrosine kinase-3 (FLT3), and TIE-2.

Pharmacodynamic effects

Cabozantinib exhibited dose-related tumour growth inhibition, tumour regression, and/or inhibited metastasis in a broad range of preclinical tumour models.

Cardiac electrophysiology

An increase from baseline in corrected QT interval by Fridericia (QTcF) of 10 – 15 ms on Day 29 (but not on Day 1) following initiation of cabozantinib treatment (at a dose of 140 mg qd) was observed in a controlled clinical study in medullary thyroid cancer patients. This effect was not associated with a change in cardiac wave form morphology or new rhythms. No cabozantinib-treated subjects in this study had a confirmed QTcF >500 ms, nor did any cabozantinib-treated subjects in the RCC study (at a dose of 60 mg).

Clinical efficacy and safety

Clinical data in renal cell carcinoma

The safety and efficacy of CABOMETYX were evaluated in a randomized, open-label, multicenter Phase 3 study. Patients (N=658) with advanced RCC with a clear cell component who had previously received at least 1 prior VEGF receptor tyrosine kinase inhibitor (VEGFR TKI) were randomized (1:1) to receive CABOMETYX (N=330) or everolimus (N=328). Patients could have received other prior therapies, including cytokines, and antibodies targeting VEGF, the programmed death 1 (PD-1) receptor, or its ligands. Patients with treated brain metastases were allowed. Progression-free survival (PFS) was assessed by a blinded independent radiology review committee, and the primary analysis was conducted among the first 375 subjects randomized. Secondary efficacy endpoints were objective response rate (ORR) and overall survival (OS). Tumor assessments were conducted every 8 weeks for the first 12 months, then every 12 weeks thereafter.

The baseline demographic and disease characteristics were similar between the CABOMETYX and everolimus arms. The majority of the patients were male (75%), with a median age of 62 years. Seventy-one percent (71%) received only one prior VEGFR TKI; 41% of patients received sunitinib as their only prior VEGFR TKI. According to the Memorial Sloan Kettering Cancer Center criteria for prognostic risk category, 46% were favorable (0 risk factors), 42% were intermediate (1 risk factor), and 13% were poor (2 or 3 risk factors). Fifty-four percent (54%) of patients had 3 or more organs with metastatic disease, including lung (63%), lymph nodes (62%), liver (29%), and bone (22%). The median duration of treatment was 7.6 months (range 0.3 – 20.5) for patients receiving CABOMETYX and 4.4 months (range 0.21 – 18.9) for patients receiving everolimus.

A statistically significant improvement in PFS was demonstrated for CABOMETYX compared to everolimus (Figure 1 and Table 3). A planned interim analysis of OS was conducted at the time of the PFS analysis and did not reach the interim boundary for statistical significance (202 events, HR=0.68 [0.51, 0.90], p=0.006). In a subsequent unplanned interim analysis of OS, a statistically significant improvement was demonstrated for

patients randomized to CABOMETYX as compared with everolimus (320 events, median of 21.4 months vs. 16.5 months; HR=0.66 [0.53, 0.83], p=0.0003; Figure 2). Comparable results for OS were observed with a follow-up analysis (descriptive) at 430 events.

Exploratory analyses of PFS and OS in the ITT population have also shown consistent results in favour of CABOMETYX compared to everolimus across different subgroups according to age (<65 vs. ≥65, sex, MSKCC risk group (favourable, intermediate, poor), ECOG status (0 vs. 1), time from diagnosis to randomisation (<1 year vs. ≥1 year), tumour MET status (high vs. low vs. unknown), bone metastases (absence vs. presence), visceral metastases (absence vs. presence), visceral and bone metastases (absence vs. presence), number of prior VEGFR-TKIs (1 vs. ≥2), duration of first VEGFR-TKI (≤6 months vs. >6 months).

Objective response rate findings are summarized in Table 4.

Figure 1: Kaplan Meier curve for progression-free survival by independent radiology review committee (first 375 randomized)





Table 3: Summary of PFS findings by independent radiology review committee

Endpoint	Primary PFS analysis Population		Intent-To-Treat Population	
	CABOMETYX	Everolimus	CABOMETYX	Everolimus
	N = 187	N = 188	N = 330	N = 328
Median PFS (95% CI), months	7.4 (5.6, 9.1)	3.8 (3.7, 5.4)	7.4 (6.6, 9.1)	3.9 (3.7, 5.1)
HR (95% CI), p-value ¹	0.58 (0.45, 0.74), p<0.0001		0.51 (0.41, 0.62), p<0.0001	

¹ stratified log-rank test

Figure 2: Kaplan-Meier curve of overall survival**Table 4: Summary of ORR findings per independent radiology committee review (IRC) and investigator review**

Endpoint	Primary Analysis ORR Intent-To-Treat Population (IRC)		ORR per Investigator Review Intent-To-Treat Population	
	CABOMETYX	Everolimus	CABOMETYX	Everolimus
	N = 330	N = 328	N = 330	N = 328
ORR (partial responses only) (95% CI)	17% (13%, 22%)	3% (2%, 6%)	24% (19%, 29%)	4% (2%, 7%)
p-value ¹	p<0.0001		p<0.0001	
Partial Response	17%	3%	24%	4%
Median time to First Response, months (95% CI)	1.91 (1.6, 11.0)	2.14 (1.9, 9.2)	1.91 (1.3, 9.8)	3.50 (1.8, 5.6)
Stable Disease as Best Response	65%	62%	63%	63%
Progressive Disease as Best Response	12%	27%	9%	27%

¹ chi-squared test

Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with CABOMETYX in all subsets of the paediatric population in treatment of kidney and renal pelvis carcinoma (excluding nephroblastoma, nephroblastomatosis, clear cell sarcoma, mesoblastic nephroma, renal medullary carcinoma and rhabdoid tumour of the kidney) (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

Absorption

Following oral administration of cabozantinib, peak cabozantinib plasma concentrations are reached at 2 to 3 hours post-dose. Plasma-concentration time profiles show a second absorption peak approximately 24 hours after administration, which suggests that cabozantinib may undergo enterohepatic recirculation.

Repeat daily dosing of cabozantinib at 140 mg for 19 days resulted in an approximately a 4- to 5-fold mean cabozantinib accumulation (based on AUC) compared to a single dose administration; steady state is achieved by approximately Day 15.

A high-fat meal moderately increased C_{max} and AUC values (41% and 57%, respectively) relative to fasted conditions in healthy volunteers administered a single 140 mg oral cabozantinib dose. There is no information on the precise food-effect when taken 1 hour after administration of cabozantinib.

Bioequivalence could not be demonstrated between the cabozantinib capsule and tablet formulations following a single 140 mg dose in healthy subjects. A 19% increase in the C_{max} of the tablet formulation (CABOMETYX) compared to the capsule formulation (COMETRIQ) was observed. A less than 10% difference in the AUC was observed between cabozantinib tablet (CABOMETYX) and capsule (COMETRIQ) formulations.

Distribution

Cabozantinib is highly protein bound *in vitro* in human plasma ($\geq 99.7\%$). Based on the population-pharmacokinetic (PK) model, the volume of distribution (V_z) is approximately 319 L (SE: $\pm 2.7\%$). Protein binding was not altered in subjects with mild or moderately impaired renal or hepatic function.

Biotransformation

Cabozantinib was metabolized *in vivo*. Four metabolites were present in plasma at exposures (AUC) greater than 10% of parent: XL184-N-oxide, XL184 amide cleavage product, XL184 monohydroxy sulfate, and 6-desmethyl amide cleavage product sulfate. Two non-conjugated metabolites (XL184-N-oxide and XL184 amide cleavage product), which possess $<1\%$ of the on-target kinase inhibition potency of parent cabozantinib, each represent $<10\%$ of total drug-related plasma exposure.

Cabozantinib is a substrate for CYP3A4 metabolism *in vitro*, as a neutralizing antibody to CYP3A4 inhibited formation of metabolite XL184 N-oxide by $>80\%$ in a NADPH-catalyzed human liver microsomal (HLM) incubation; in contrast, neutralizing antibodies to CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C19, CYP2D6 and CYP2E1 had no effect on cabozantinib metabolite formation. A neutralizing antibody to CYP2C9 showed a minimal effect on cabozantinib metabolite formation (ie, a $<20\%$ reduction).

Elimination

In a population PK analysis of cabozantinib using data collected from 318 patients with RCC and 63 normal healthy volunteers following oral administration of doses of 60 mg, 40 mg, and 20 mg, the plasma terminal half-life of cabozantinib is approximately 99 hours. Mean clearance (CL/F) at steady-state was estimated to be 2.2 L/hr. Within a 48-day collection period after a single dose of ^{14}C -cabozantinib in healthy volunteers, approximately 81% of the total administered radioactivity was recovered with 54% in faeces and 27% in urine.

Pharmacokinetics in special patient populations

Renal impairment

Results from a study in patients with renal impairment indicate that the ratios of geometric LS mean for plasma cabozantinib, C_{max} and AUC_{0-inf} were 19% and 30% higher, for subjects with mild renal impairment (90% CI for C_{max} 91.60% to 155.51%; AUC_{0-inf} 98.79% to 171.26%) and 2% and 6-7% higher (90% CI for C_{max} 78.64% to 133.52%; AUC_{0-inf} 79.61% to 140.11%), for subjects with moderate renal impairment

compared to subjects with normal renal function. Patients with severe renal impairment have not been studied.

Hepatic impairment

Results from a study in patients with hepatic impairment indicate that exposure (AUC_{0-inf}) increased by 81% and 63% in subjects with mild and moderate hepatic impairment, respectively (90% CI for AUC_{0-inf} : 121.44% to 270.34% for mild and 107.37% to 246.67% for moderate). Patients with severe hepatic impairment have not been studied.

Race

A population PK analysis did not identify clinically relevant differences in PK of cabozantinib based on race.

5.3 Preclinical safety data

Adverse reactions not observed in clinical studies, but seen in animals at exposure levels similar to clinical exposure levels and with possible relevance to clinical use were as follows:

In rat and dog repeat-dose toxicity studies up to 6 months duration, target organs for toxicity were GI tract, bone marrow, lymphoid tissues, kidney, adrenal and reproductive tract tissues. The no observed adverse effect level (NOAEL) for these findings were below human clinical exposure levels at intended therapeutic dose.

Cabozantinib has shown no mutagenic or clastogenic potential in a standard battery of genotoxicity assays. The carcinogenic potential of cabozantinib has been evaluated in two species: rasH2 transgenic mice and Sprague-Dawley rats. In the 2-year rat carcinogenicity study, cabozantinib-related neoplastic findings consisted of an increased incidence of benign pheochromocytoma, alone or in combination with malignant pheochromocytoma/complex malignant pheochromocytoma of the adrenal medulla in both sexes at exposures well below the intended exposure in humans. The clinical relevance of the observed neoplastic lesions in rats is uncertain, but likely to be low.

Cabozantinib was not carcinogenic in the rasH2 mouse model at a slightly higher exposure than the intended human therapeutic exposure.

Fertility studies in rats have shown reduced male and female fertility. Further, hypospermatogenesis was observed in male dogs at exposure levels below human clinical exposure levels at intended therapeutic dose.

Embryo-foetal development studies were performed in rats and rabbits. In rats, cabozantinib caused postimplantation loss, foetal oedema, cleft palate/lip, dermal aplasia and kinked or rudimentary tail. In rabbits, cabozantinib produced foetal soft tissue changes (reduced spleen size, small or missing intermediate lung lobe) and increased foetal incidence of total malformations. NOAEL for embryo-foetal toxicity and teratogenic findings were below human clinical exposure levels at intended therapeutic dose.

Juvenile rats (comparable to a >2 year old pediatric population) administered cabozantinib showed increased WBC parameters, decreased haematopoiesis, pubescent/immature female reproductive system (without delayed vaginal opening), tooth abnormalities, reduced bone mineral content and density, liver pigmentation and lymph node lymphoid hyperplasia. Findings in uterus/ovaries and decreased haematopoiesis appeared to be transient, while effects on bone parameters and liver pigmentation were sustained. Juvenile rats (correlating to a <2 year pediatric population) showed similar treatment-related findings, but appeared to be more sensitive to cabozantinib-related toxicity at comparable dose levels.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet content

Microcrystalline cellulose
Anhydrous lactose
Hydroxypropyl cellulose
Croscarmellose sodium
Colloidal anhydrous silica
Magnesium stearate

Film-coating

Hypromellose 2910
Titanium dioxide (E171)
Triacetin
Iron oxide yellow (E172)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

PVC/PCTFE blister with push-through aluminum foil backing containing 7 film-coated tablets. Each carton contains 4 blisters with 28 film-coated tablets.

HDPE bottle with a polypropylene child-resistant closure and three silica gel dessicant canisters. Each bottle contains 30 film-coated tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Ipsen Pharma
65 quai Georges Gorse
92100 Boulogne-Billancourt
France

8. MARKETING AUTHORISATION NUMBER(S)

Cabometyx 20 mg film-coated tablets

EU/1/16/1136/001

EU/1/16/1136/002

Cabometyx 40 mg film-coated tablets

EU/1/16/1136/003

EU/1/16/1136/004

Cabometyx 60 mg film-coated tablets

EU/1/16/1136/005

EU/1/16/1136/006

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

ANNEX II

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer(s) responsible for batch release

Patheon France
40 Boulevard de Champaret
38300 Bourgoin-Jallieu
FRANCE

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription.

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

•Periodic safety update reports

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder shall submit the first periodic safety update report for this product within 6 months following authorisation.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

•Risk Management Plan (RMP)

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

OUTER CARTON

1. NAME OF THE MEDICINAL PRODUCT

CABOMETYX 20 mg film-coated tablets
cabozantinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each tablet contains cabozantinib (*S*)-malate equivalent to 20 mg of cabozantinib.

3. LIST OF EXCIPIENTS

Contains lactose. See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet
28 film-coated tablets
30 film-coated tablets

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

Any unused product or waste material should be disposed of in accordance with local requirements.

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Ipsen Pharma
65 quai Georges Gorse
92100 Boulogne-Billancourt
France

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1136/001
EU/1/16/1136/002

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

CABOMETYX 20 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

CABOMETYX 20 mg film-coated tablets
cabozantinib

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Ipsen Pharma

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

OUTER CARTON

1. NAME OF THE MEDICINAL PRODUCT

CABOMETYX 40 mg film-coated tablets
cabozantinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each tablet contains cabozantinib (*S*)-malate equivalent to 40 mg of cabozantinib.

3. LIST OF EXCIPIENTS

Contains lactose. See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet
28 film-coated tablets
30 film-coated tablets

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

Any unused product or waste material should be disposed of in accordance with local requirements.

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Ipsen Pharma
65 quai Georges Gorse
92100 Boulogne-Billancourt
France

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1136/003
EU/1/16/1136/004

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

CABOMETYX 40 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

CABOMETYX 40 mg film-coated tablets
cabozantinib

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Ipsen Pharma

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

OUTER CARTON

1. NAME OF THE MEDICINAL PRODUCT

CABOMETYX 60 mg film-coated tablets
cabozantinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each tablet contains cabozantinib (S)-malate equivalent to 60 mg of cabozantinib.

3. LIST OF EXCIPIENTS

Contains lactose. See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet
28 film-coated tablets
30 film-coated tablets

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

Any unused product or waste material should be disposed of in accordance with local requirements.

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Ipsen Pharma
65 quai Georges Gorse
92100 Boulogne-Billancourt
France

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1136/005
EU/1/16/1136/006

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

CABOMETYX 60 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

CABOMETYX 60 mg film-coated tablets
cabozantinib

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Ipsen Pharma

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING

BOTTLE LABEL

1. NAME OF THE MEDICINAL PRODUCT

CABOMETYX 20 mg film-coated tablets
cabozantinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each tablet contains cabozantinib (*S*)-malate equivalent to 20 mg cabozantinib.

3. LIST OF EXCIPIENTS

Contains lactose. See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

30 film-coated tablets

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Ipsen Pharma
65 quai Georges Gorse
92100 Boulogne-Billancourt
France

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1136/002

13. BATCH NUMBER<, DONATION AND PRODUCT CODES>

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING

BOTTLE LABEL

1. NAME OF THE MEDICINAL PRODUCT

CABOMETYX 40 mg film-coated tablets
cabozantinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each tablet contains cabozantinib (*S*)-malate equivalent to 40 mg cabozantinib.

3. LIST OF EXCIPIENTS

Contains lactose. See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

30 film-coated tablets

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Ipsen Pharma
65 quai Georges Gorse
92100 Boulogne-Billancourt
France

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1136/004

13. BATCH NUMBER<, DONATION AND PRODUCT CODES>

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING

BOTTLE LABEL

1. NAME OF THE MEDICINAL PRODUCT

CABOMETYX 60 mg film-coated tablets
cabozantinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each tablet contains cabozantinib (*S*)-malate equivalent to 60 mg cabozantinib.

3. LIST OF EXCIPIENTS

Contains lactose. See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

30 film-coated tablets

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Ipsen Pharma
65 quai Georges Gorse
92100 Boulogne-Billancourt
France

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1136/006

13. BATCH NUMBER<, DONATION AND PRODUCT CODES>

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

B. PACKAGE LEAFLET

Package leaflet: Information for the patient

CABOMETYX 20 mg film-coated tablets
CABOMETYX 40 mg film-coated tablets
CABOMETYX 60 mg film-coated tablets
cabozantinib

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What CABOMETYX is and what it is used for
2. What you need to know before you take CABOMETYX
3. How to take CABOMETYX
4. Possible side effects
5. How to store CABOMETYX
6. Contents of the pack and other information

1. What CABOMETYX is and what it is used for

What CABOMETYX is

CABOMETYX is a cancer medicine that contains the active substance cabozantinib. It is used to treat advanced stages of a type of kidney cancer called renal cell carcinoma, in adults who have already had previous treatment.

How CABOMETYX works

CABOMETYX blocks the action of proteins called receptor tyrosine kinases (RTKs), which are involved in the growth of cells and the development of new blood vessels that supply them. These proteins can be present in high amounts in cancer cells, and by blocking their action CABOMETYX can slow down the rate at which the tumour grows and help to cut off the blood supply that the cancer needs.

2. What you need to know before you take CABOMETYX

Do not take CABOMETYX

- if you are allergic to cabozantinib or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor or pharmacist before taking CABOMETYX if you:

- have high blood pressure
- have diarrhoea
- have a recent history of significant bleeding
- have had surgery within the last month (or if surgical procedures are planned), including dental surgery
- have inflammatory bowel disease (for example, Crohn's disease or ulcerative colitis, diverticulitis, or appendicitis)
- have a recent history of blood clot in the leg, stroke, or heart attack
- have liver or kidney disease.

Tell your doctor if any of these affect you.

You may need treatment for them, or your doctor may decide to change your dose of CABOMETYX, or stop treatment altogether. See also section 4 "*Possible side effects*".

Children and adolescents

CABOMETYX is not recommended for children or adolescents. The effects of CABOMETYX in people younger than 18 years old are not known.

Other medicines and CABOMETYX

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. This is because CABOMETYX can affect the way some other medicines work. Also, some medicines can affect the way CABOMETYX works. This could mean that your doctor needs to change the dose(s) that you take. You should tell your doctor about every medicines, but in particular if taking:

- Medicines that treat fungal infections, such as itraconazole, ketoconazole, and posaconazole
- Medicines used to treat bacterial infections (antibiotics) such as erythromycin, clarithromycin, and rifampicin
- Allergy medicines such as fexofenadine and ranolazine
- Medicines used to treat epilepsy or fits such as phenytoin, carbamazepine, and phenobarbital
- Herbal preparations containing St. John's Wort (*Hypericum perforatum*), sometimes used for treating depression or depression-related conditions such as anxiety
- Medicines used to thin the blood, such as warfarin
- Medicines to treat high blood pressure or other heart conditions, such as aliskiren, ambrisentan, dabigatran etexilate, digoxin, talinolol, and tolvaptan
- Medicines for diabetes, such as saxagliptin and sitagliptin
- Medicines used to treat gout, such as colchicine
- Medicines used to treat HIV or AIDS, such as efavirenz, ritonavir, maraviroc and emtricitabine
- Medicines used to prevent transplant rejection (ciclosporin) and ciclosporin-based regimens in rheumatoid arthritis and psoriasis

Oral contraceptives

If you take CABOMETYX whilst using oral contraceptives, the oral contraceptives may be ineffective. You should also use a barrier contraceptive (e.g. condom or diaphragm) whilst taking CABOMETYX and for at least 4 months after treatment has finished.

Taking CABOMETYX with food

You should not take CABOMETYX with food. You should not eat anything for at least 2 hours before taking CABOMETYX and for 1 hour after taking the medicine. Avoid consuming grapefruit-containing products for as long as you are using this medicine, as they may increase the levels of CABOMETYX in your blood.

Pregnancy, breast-feeding and fertility

Avoid becoming pregnant while being treated with CABOMETYX. If you or your partner could become pregnant, use adequate contraception during treatment and for at least 4 months after treatment has finished. Talk to your doctor about which methods of contraception are appropriate while you are taking CABOMETYX (see also under Other medicines and CABOMETYX, above).

Tell your doctor if you or your partner become pregnant or plan to become pregnant while you are being treated with CABOMETYX.

Talk to your doctor BEFORE taking CABOMETYX if you or your partner are considering or planning to have a baby after your treatment has finished. There is a possibility your fertility could be affected by treatment with CABOMETYX.

Women taking CABOMETYX should not breast-feed during treatment and for at least 4 months after treatment has finished, as cabozantinib and/or its metabolites may be excreted in breast milk and be harmful to your child.

Driving and using machines

Use caution when driving or using machines. Keep in mind that treatment with CABOMETYX may make you feel tired or weak and can affect your ability to drive or use machines.

CABOMETYX contains lactose

CABOMETYX contains lactose (a type of sugar). If you have been told by your doctor that you have an intolerance to some sugars, talk to your doctor before taking this medicine.

3. How to take CABOMETYX

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

You should continue to take this medicine until your doctor decides to stop your treatment. If you get serious side effects, your doctor may decide to change your dose or stop treatment earlier than originally planned. Your doctor will tell you if you need your dose adjusted.

CABOMETYX should be taken once a day. The usual dose is 60 mg, however your doctor will decide on the right dose for you.

CABOMETYX should **not** be taken with food. You should not eat anything for at least 2 hours before taking CABOMETYX and for 1 hour after taking the medicine. Swallow the tablet with a full glass of water. Do not crush the tablets.

If you take more CABOMETYX than you should

If you have taken more CABOMETYX than you have been instructed to, talk to a doctor or go to the hospital with the tablets and this leaflet straight away.

If you forget to take CABOMETYX

- If there are still 12 hours or more before your next dose is due, then take the missed dose as soon as you remember. Take the next dose at the normal time.
- If your next dose is due in less than 12 hours, then do not take the dose that you have missed. Take your next dose at the normal time.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. If you get side effects, your doctor may tell you to take CABOMETYX at a lower dose. Your doctor may also prescribe other medicines to help control your side effects.

Tell your doctor straight away if you notice any of the following side effects – you may need urgent medical treatment:

- Symptoms including pain in the abdomen (belly), nausea (feeling sick), vomiting, constipation, or fever. These may be signs of a gastrointestinal perforation, a hole that develops in your stomach or intestine that could be life-threatening.
- Severe or uncontrollable bleeding with symptoms such as: vomiting blood, black stools, bloody urine, headache, coughing up of blood.
- Swelling, pain in your hands and feet, or shortness of breath.
- A wound that does not heal.
- Fits, headaches, confusion, or finding it difficult to concentrate. These may be signs of a condition called reversible posterior leukoencephalopathy syndrome (RPLS). RPLS is uncommon (it affects less than 1 in 100 people).

Other side effects include:

Very common side effects (may affect more than 1 in 10 people)

- Stomach upset, including diarrhoea, nausea, vomiting, constipation, indigestion, and abdominal pain
- Blisters, pain of the hands or soles of the feet, rash or redness of the skin, dry skin
- Decreased appetite, weight loss, altered sense of taste
- Fatigue, weakness, headache, dizziness
- Hypertension (increase in blood pressure)
- Anaemia (low levels of red blood cells)
- Redness, swelling or pain in the mouth or throat, difficulty in speaking, hoarseness, cough
- Changes in blood tests used to monitor general health and function of your organs (including the liver), low levels of electrolytes (like magnesium, calcium, sodium, or potassium)
- Increase in the level of bilirubin in your blood (which may result in jaundice/yellow skin or eyes)
- Pain in arms, legs and joints, muscle spasms
- Shortness of breath
- Protein in urine (seen in tests)
- Reduced thyroid activity; symptoms can include: tiredness, weight gain, constipation, feeling cold and dry skin

Common side effects (may affect up to 1 in 10 people)

- Abscess (collection of pus, with swelling and inflammation)
- Dehydration (lack of fluids)
- Ringing in ears
- Blood clots in the lungs

- Pain in the upper part of the abdomen
- Gastro-oesophageal reflux disease (bringing up stomach acid)
- Haemorrhoids (piles)
- Alopecia (hair loss and thinning)
- Swelling in your legs, feet, arms and hands

Uncommon side effects (may affect 1 in 100 people)

- Fits
- A painful tear or abnormal connection of the tissue in your anus
- Inflammation of the pancreas
- Decrease in bile flow from the liver
- Bone damage in the jaw

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store CABOMETYX

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister, bottle label and carton after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What CABOMETYX contains

The active substance is cabozantinib (*S*)-malate.

CABOMETYX 20 mg film-coated tablets: Each tablet contains cabozantinib (*S*)-malate equivalent to 20 mg of cabozantinib.

CABOMETYX 40 mg film-coated tablets: Each tablet contains cabozantinib (*S*)-malate equivalent to 40 mg of cabozantinib.

CABOMETYX 60 mg film-coated tablets: Each tablet contains cabozantinib (*S*)-malate equivalent to 60 mg of cabozantinib.

The other ingredients are:

- **Tablet contents:** microcrystalline cellulose, lactose anhydrous, hydroxypropyl cellulose, croscarmellose sodium, colloidal silicon dioxide anhydrous, magnesium stearate. (see section 2 for lactose content)
- **Film coating:** hypromellose, titanium dioxide (E171), triacetin, iron oxide yellow (E172)

What CABOMETYX looks like and contents of the pack

CABOMETYX 20 mg film-coated tablets are yellow, round with no score, and identified with “XL” on one side and “20” on the other side.

CABOMETYX 40 mg film-coated tablets are yellow, triangle shaped with no score, and identified with “XL” on one side and “40” on the other side.

CABOMETYX 60 mg film-coated tablets are yellow, oval shaped with no score, and identified with “XL” on one side and “60” on the other side.

CABOMETYX tablets are available in packs containing either 4 blisters with 7 tablets each (28 total), or one plastic bottle with 30 tablets.

Not all pack sizes may be marketed in your country.

Marketing Authorisation Holder

Ipsen Pharma
65 quai Georges Gorse
92100 Boulogne-Billancourt
France

Manufacturer

Patheon France
40 Boulevard de Champaret
38300 Bourgoin Jallieu, France

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

België/Belgique/Belgien, Luxembourg/Luxemburg

Ipsen NV Guldensporenpark 87
B-9820 Merelbeke
België /Belgique/Belgien
Tél/Tel: + 32 - 9 - 243 96 00

Italia

Ipsen SpA
Via del Bosco Rinnovato n. 6
Milanofiori Nord Palazzo U7
20090 Assago (Mi)
Tel: + 39 - 02 - 39 22 41

България

PharmaSwiss EOOD
16, Troyanski Prohod Street,
Floor 3, Office 8, Lagera
1612 Sofia
Тел.: +359 2 8952 110

Latvija

Ipsen Pharma representative office
Kalnciema street 33-5
Riga
LV 1046
Tel: +371 67622233

Česká republika

Ipsen Pharma, o.s. Olbrachtova 2006/9,
140 00 Praha 4
Tel: + 420 242 481 821

Lietuva

Ipsen Pharma SAS Lietuvos filialas
Narbuto g. 5,
08103 Vilnius
Tel. +370 700 33305

Danmark, Norge, Suomi/Finland, Sverige, Ísland

Institut Produits Synthèse (IPSEN) AB
Kista Science Tower
Färögatan 33

Magyarország

Ipsen Pharma SAS Magyarországi Kereskedelmi
Képviselő
Árbóc utca 6.

SE- 164 51 Kista
Sverige/Ruotsi/Svíþjóð
Tlf/Puh/Tel/Sími: +46 8 451 60 00

Deutschland, Österreich

Ipsen Pharma GmbH
Willy-Brandt-Str. 3
D-76275 Ettlingen
Tel.: +49 7243 184-80

Eesti

ESTOBIIN OÜ
Udeselja 4-4
EE-11913 Tallinn
Tel: +372 51 55 810

Ελλάδα, Κύπρος, Malta

Ipsen ΕΠΕ
Αγ. Δημητρίου 63 Άλιμος
GR-17456 Αθήνα Ελλάδα
Τηλ: + 30 - 210 - 984 3324

España

Ipsen Pharma, S.A.
Torre Realía, Plaza de Europa, 41-43
08908 L'Hospitalet de Llobregat
Barcelona
Tel: + 34 - 936 - 858 100

France

Ipsen Pharma
65 quai Georges Gorse
92100 Boulogne-Billancourt
France
Tél: + 33 1 58 33 50 00

Hrvatska

PharmaSwiss d.o.o.
Strojarska 20, 10 000 Zagreb,
Hrvatska
Tel: +385 1 6311 833
Fax: +385 1 6311 844

Ireland

Ipsen Pharmaceuticals Ltd.
Blanchardstown Industrial Park
Blanchardstown
IRL-Dublin 15
Tel: +353-1-809-8256

H- 1133 Budapest
Tel.: +36-1-555-5930

Nederland

Ipsen Farmaceutica B.V.
Taurusavenue 33b
2132 LS Hoofddorp
Tel: + 31 (0) 23 554 1600

Polska

Ipsen Poland Sp. z o.o. Al. Jana Pawła II 29
00-867 Warszawa
Tel.: + 48 (0) 22 653 68 00

Portugal

Ipsen Portugal - Produtos Farmacêuticos S.A.
Alameda Fernão Lopes, n° 16-11°
Miraflores P-1495 - 190 Algés
Portugal
Tel: + 351 - 21 - 412 3550

România

Ipsen Pharma
Str. Grigore Alexandrescu nr. 59,
clădirea HQ Victoriei, Sector 1,
010626, București
Tel: + 40 (021) 231 27 20

Slovenija

PharmaSwiss d.o.o.
Brodišče 32
SI-1236 Trzin
Tel: + 386 1 236 47 00

Slovenská republika

Liek s.r.o.
Hviezdoslavova 19
SK-90301 Senec
Slovenská republika
Tel: + 421 253 412 018

United Kingdom

Ipsen Ltd.
190 Bath Road
Slough, Berkshire SL1 3XE
United Kingdom
Tel: + 44 (0)1753 - 62 77 00

This leaflet was last revised in

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu>.

ANNEX IV

**CONCLUSIONS ON THE REQUEST FOR ONE-YEAR DATA EXCLUSIVITY PRESENTED BY
THE EUROPEAN MEDICINES AGENCY**

Conclusions presented by the European Medicines Agency on:

•one-year marketing protection

The CHMP reviewed the data submitted by the marketing authorisation holder, taking into account the provisions of Article 14(11) of Regulation (EC) No 726/2004, and considers that the new therapeutic indication brings significant clinical benefit in comparison with existing therapies as further explained in the European Public Assessment Report.