

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## 1. NAME OF THE MEDICINAL PRODUCT

Flixabi 100 mg powder for concentrate for solution for infusion.

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One vial contains 100 mg of infliximab\*. After reconstitution, each ml contains 10 mg of infliximab.

\* Infliximab is a chimeric human-murine IgG1 monoclonal antibody produced in Chinese Hamster Ovary (CHO) cells by recombinant DNA technology.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Powder for concentrate for solution for infusion

White powder.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

#### Rheumatoid arthritis

Flixabi, in combination with methotrexate, is indicated for the reduction of signs and symptoms as well as the improvement in physical function in:

- adult patients with active disease when the response to disease-modifying antirheumatic drugs (DMARDs), including methotrexate, has been inadequate.
- adult patients with severe, active and progressive disease not previously treated with methotrexate or other DMARDs.

In these patient populations, a reduction in the rate of the progression of joint damage, as measured by X-ray, has been demonstrated (see section 5.1).

#### Adult Crohn's disease

Flixabi is indicated for:

- treatment of moderately to severely active Crohn's disease, in adult patients who have not responded despite a full and adequate course of therapy with a corticosteroid and/or an immunosuppressant; or who are intolerant to or have medical contraindications for such therapies.
- treatment of fistulising, active Crohn's disease, in adult patients who have not responded despite a full and adequate course of therapy with conventional treatment (including antibiotics, drainage and immunosuppressive therapy).

#### Paediatric Crohn's disease

Flixabi is indicated for treatment of severe, active Crohn's disease in children and adolescents aged 6 to 17 years, who have not responded to conventional therapy including a corticosteroid, an immunomodulator and primary nutrition therapy; or who are intolerant to or have contraindications for such therapies. Infliximab has been studied only in combination with conventional immunosuppressive therapy.

### Ulcerative colitis

Flixabi is indicated for treatment of moderately to severely active ulcerative colitis in adult patients who have had an inadequate response to conventional therapy including corticosteroids and 6-mercaptopurine (6-MP) or azathioprine (AZA), or who are intolerant to or have medical contraindications for such therapies.

### Paediatric ulcerative colitis

Flixabi is indicated for treatment of severely active ulcerative colitis in children and adolescents aged 6 to 17 years, who have had an inadequate response to conventional therapy including corticosteroids and 6-MP or AZA, or who are intolerant to or have medical contraindications for such therapies.

### Ankylosing spondylitis

Flixabi is indicated for treatment of severe, active ankylosing spondylitis, in adult patients who have responded inadequately to conventional therapy.

### Psoriatic arthritis

Flixabi is indicated for treatment of active and progressive psoriatic arthritis in adult patients when the response to previous DMARD therapy has been inadequate.

Flixabi should be administered

- in combination with methotrexate
- or alone in patients who show intolerance to methotrexate or for whom methotrexate is contraindicated.

Infliximab has been shown to improve physical function in patients with psoriatic arthritis, and to reduce the rate of progression of peripheral joint damage as measured by X-ray in patients with polyarticular symmetrical subtypes of the disease (see section 5.1).

### Psoriasis

Flixabi is indicated for treatment of moderate to severe plaque psoriasis in adult patients who failed to respond to, or who have a contraindication to, or are intolerant to other systemic therapy including cyclosporine, methotrexate or psoralen ultra-violet A (PUVA) (see section 5.1).

## **4.2 Posology and method of administration**

Flixabi treatment is to be initiated and supervised by qualified physicians experienced in the diagnosis and treatment of rheumatoid arthritis, inflammatory bowel diseases, ankylosing spondylitis, psoriatic arthritis or psoriasis. Flixabi should be administered intravenously. Flixabi infusions should be administered by qualified healthcare professionals trained to detect any infusion-related issues. Patients treated with Flixabi should be given the package leaflet and the special Alert card.

During Flixabi treatment, other concomitant therapies, e.g. corticosteroids and immunosuppressants should be optimised.

### Posology

#### *Adults (≥ 18 years)*

#### Rheumatoid arthritis

3 mg/kg given as an intravenous infusion followed by additional 3 mg/kg infusion doses at 2 and 6 weeks after the first infusion, then every 8 weeks thereafter.

Flixabi must be given concomitantly with methotrexate.

Available data suggest that the clinical response is usually achieved within 12 weeks of treatment. If a patient has an inadequate response or loses response after this period, consideration may be given to increase the dose step-wise by approximately 1.5 mg/kg, up to a maximum of 7.5 mg/kg every 8 weeks.

Alternatively, administration of 3 mg/kg as often as every 4 weeks may be considered. If adequate response is achieved, patients should be continued on the selected dose or dose frequency. Continued therapy should

be carefully reconsidered in patients who show no evidence of therapeutic benefit within the first 12 weeks of treatment or after dose adjustment.

#### Moderately to severely active Crohn's disease

5 mg/kg given as an intravenous infusion followed by an additional 5 mg/kg infusion 2 weeks after the first infusion. If a patient does not respond after 2 doses, no additional treatment with Flixabi should be given. Available data do not support further Flixabi treatment, in patients not responding within 6 weeks of the initial infusion.

In responding patients, the alternative strategies for continued treatment are:

- Maintenance: Additional infusion of 5 mg/kg at 6 weeks after the initial dose, followed by infusions every 8 weeks or
- Re-administration: Infusion of 5 mg/kg if signs and symptoms of the disease recur (see 'Re-administration' below and section 4.4).

Although comparative data are lacking, limited data in patients who initially responded to 5 mg/kg but who lost response indicate that some patients may regain response with dose escalation (see section 5.1). Continued therapy should be carefully reconsidered in patients who show no evidence of therapeutic benefit after dose adjustment.

#### Fistulising, active Crohn's disease

5 mg/kg given as an intravenous infusion followed by additional 5 mg/kg infusions at 2 and 6 weeks after the first infusion. If a patient does not respond after 3 doses, no additional treatment with Flixabi should be given.

In responding patients, the alternative strategies for continued treatment are:

- Maintenance: Additional infusions of 5 mg/kg every 8 weeks or
- Re-administration: Infusion of 5 mg/kg if signs and symptoms of the disease recur followed by infusions of 5 mg/kg every 8 weeks (see 'Re-administration' below and section 4.4).

Although comparative data are lacking, limited data in patients who initially responded to 5 mg/kg but who lost response indicate that some patients may regain response with dose escalation (see section 5.1). Continued therapy should be carefully reconsidered in patients who show no evidence of therapeutic benefit after dose adjustment.

In Crohn's disease, experience with re-administration if signs and symptoms of disease recur is limited and comparative data on the benefit/risk of the alternative strategies for continued treatment are lacking.

#### Ulcerative colitis

5 mg/kg given as an intravenous infusion followed by additional 5 mg/kg infusion doses at 2 and 6 weeks after the first infusion, then every 8 weeks thereafter.

Available data suggest that the clinical response is usually achieved within 14 weeks of treatment, i.e. three doses. Continued therapy should be carefully reconsidered in patients who show no evidence of therapeutic benefit within this time period.

#### Ankylosing spondylitis

5 mg/kg given as an intravenous infusion followed by additional 5 mg/kg infusion doses at 2 and 6 weeks after the first infusion, then every 6 to 8 weeks. If a patient does not respond by 6 weeks (i.e. after 2 doses), no additional treatment with Flixabi should be given.

#### Psoriatic arthritis

5 mg/kg given as an intravenous infusion followed by additional 5 mg/kg infusion doses at 2 and 6 weeks after the first infusion, then every 8 weeks thereafter.

#### Psoriasis

5 mg/kg given as an intravenous infusion followed by additional 5 mg/kg infusion doses at 2 and 6 weeks after the first infusion, then every 8 weeks thereafter. If a patient shows no response after 14 weeks (i.e. after 4 doses), no additional treatment with Flixabi should be given.

#### Re-administration for Crohn's disease and rheumatoid arthritis

If the signs and symptoms of disease recur, Flixabi can be re-administered within 16 weeks following the last infusion. In clinical studies, delayed hypersensitivity reactions have been uncommon and have occurred after infliximab-free intervals of less than 1 year (see sections 4.4 and 4.8). The safety and efficacy of re-administration after an infliximab-free interval of more than 16 weeks has not been established. This applies to both Crohn's disease patients and rheumatoid arthritis patients.

#### Re-administration for ulcerative colitis

The safety and efficacy of re-administration, other than every 8 weeks, has not been established (see sections 4.4 and 4.8).

#### Re-administration for ankylosing spondylitis

The safety and efficacy of re-administration, other than every 6 to 8 weeks, has not been established (see sections 4.4 and 4.8).

#### Re-administration for psoriatic arthritis

The safety and efficacy of re-administration, other than every 8 weeks, has not been established (see sections 4.4 and 4.8).

#### Re-administration for psoriasis

Limited experience from re-treatment with one single infliximab dose in psoriasis after an interval of 20 weeks suggests reduced efficacy and a higher incidence of mild to moderate infusion reactions when compared to the initial induction regimen (see section 5.1).

Limited experience from re-treatment following disease flare by a re-induction regimen suggests a higher incidence of infusion reactions, including serious ones, when compared to 8-weekly maintenance treatment (see section 4.8).

#### Re-administration across indications

In case maintenance therapy is interrupted, and there is a need to restart treatment, use of a re-induction regimen is not recommended (see section 4.8). In this situation, Flixabi should be re-initiated as a single dose followed by the maintenance dose recommendations described above.

#### *Elderly ( $\geq 65$ years)*

Specific studies of infliximab in elderly patients have not been conducted. No major age-related differences in clearance or volume of distribution were observed in clinical studies. No dose adjustment is required (see section 5.2). For more information about the safety of infliximab in elderly patients see sections 4.4 and 4.8.

#### *Renal and/or hepatic impairment*

Infliximab has not been studied in these patient populations. No dose recommendations can be made (see section 5.2).

#### *Paediatric population*

##### Crohn's disease (6 to 17 years)

5 mg/kg given as an intravenous infusion followed by additional 5 mg/kg infusion doses at 2 and 6 weeks after the first infusion, then every 8 weeks thereafter. Available data do not support further infliximab treatment in children and adolescents not responding within the first 10 weeks of treatment (see section 5.1).

Some patients may require a shorter dosing interval to maintain clinical benefit, while for others a longer dosing interval may be sufficient. Patients who have had their dose interval shortened to less than 8 weeks

may be at greater risk for adverse reactions. Continued therapy with a shortened interval should be carefully considered in those patients who show no evidence of additional therapeutic benefit after a change in dosing interval.

The safety and efficacy of infliximab have not been studied in children with Crohn's disease below the age of 6 years. Currently available pharmacokinetic data are described in section 5.2 but no recommendation on a posology can be made in children younger than 6 years.

#### Ulcerative colitis (6 to 17 years)

5 mg/kg given as an intravenous infusion followed by additional 5 mg/kg infusion doses at 2 and 6 weeks after the first infusion, then every 8 weeks thereafter. Available data do not support further infliximab treatment in paediatric patients not responding within the first 8 weeks of treatment (see section 5.1).

The safety and efficacy of infliximab have not been studied in children with ulcerative colitis below the age of 6 years. Currently available pharmacokinetic data are described in section 5.2 but no recommendation on a posology can be made in children younger than 6 years.

#### Psoriasis

The safety and efficacy of infliximab in children and adolescents younger than 18 years for the indication of psoriasis have not been established. Currently available data are described in section 5.2 but no recommendation on a posology can be made.

#### Juvenile idiopathic arthritis, psoriatic arthritis and ankylosing spondylitis

The safety and efficacy of infliximab in children and adolescents younger than 18 years for the indications of juvenile idiopathic arthritis, psoriatic arthritis and ankylosing spondylitis have not been established. Currently available data are described in section 5.2 but no recommendation on a posology can be made.

#### Juvenile rheumatoid arthritis

The safety and efficacy of infliximab in children and adolescents younger than 18 years for the indication of juvenile rheumatoid arthritis have not been established. Currently available data are described in sections 4.8 and 5.2 but no recommendation on a posology can be made.

#### *Renal and/or hepatic impairment*

Infliximab has not been studied in these patient populations. No dose recommendations can be made (see section 5.2).

#### Method of administration

Flixabi should be administered intravenously over a 2-hour period. All patients administered Flixabi are to be observed for at least 1-2 hours post-infusion for acute infusion-related reactions. Emergency equipment, such as adrenaline, antihistamines, corticosteroids and an artificial airway must be available. Patients may be pre-treated with e.g., an antihistamine, hydrocortisone and/or paracetamol and infusion rate may be slowed in order to decrease the risk of infusion-related reactions especially if infusion-related reactions have occurred previously (see section 4.4).

#### *Shortened infusions across adult indications*

In carefully selected adult patients who have tolerated at least 3 initial 2-hour infusions of Flixabi (induction phase) and are receiving maintenance therapy, consideration may be given to administering subsequent infusions over a period of not less than 1 hour. If an infusion reaction occurs in association with a shortened infusion, a slower infusion rate may be considered for future infusions if treatment is to be continued. Shortened infusions at doses > 6 mg/kg have not been studied (see section 4.8).

For preparation and administration instructions, see section 6.6.

### **4.3 Contraindications**

Patients with a history of hypersensitivity to infliximab (see section 4.8), to other murine proteins, or to any of the excipients listed in section 6.1.

Patients with tuberculosis or other severe infections such as sepsis, abscesses, and opportunistic infections (see section 4.4).

Patients with moderate or severe heart failure (NYHA class III/IV) (see sections 4.4 and 4.8).

#### **4.4 Special warnings and precautions for use**

In order to improve the traceability of biological medicinal products, the trademark and the batch number of the administered product should be clearly recorded (or stated) in the patient file.

##### Infusion reactions and hypersensitivity

Infliximab has been associated with acute infusion-related reactions, including anaphylactic shock, and delayed hypersensitivity reactions (see section 4.8).

Acute infusion reactions including anaphylactic reactions may develop during (within seconds) or within a few hours following infusion. If acute infusion reactions occur, the infusion must be interrupted immediately. Emergency equipment, such as adrenaline, antihistamines, corticosteroids and an artificial airway must be available. Patients may be pre-treated with e.g., an antihistamine, hydrocortisone and/or paracetamol to prevent mild and transient effects.

Antibodies to infliximab may develop and have been associated with an increased frequency of infusion reactions. A low proportion of the infusion reactions was serious allergic reactions. An association between development of antibodies to infliximab and reduced duration of response has also been observed. Concomitant administration of immunomodulators has been associated with lower incidence of antibodies to infliximab and a reduction in the frequency of infusion reactions. The effect of concomitant immunomodulator therapy was more profound in episodically-treated patients than in patients given maintenance therapy. Patients who discontinue immunosuppressants prior to or during infliximab treatment are at greater risk of developing these antibodies. Antibodies to infliximab cannot always be detected in serum samples. If serious reactions occur, symptomatic treatment must be given and further Flixabi infusions must not be administered (see section 4.8).

In clinical studies, delayed hypersensitivity reactions have been reported. Available data suggest an increased risk for delayed hypersensitivity with increasing infliximab-free interval. Patients should be advised to seek immediate medical advice if they experience any delayed adverse event (see section 4.8). If patients are re-treated after a prolonged period, they must be closely monitored for signs and symptoms of delayed hypersensitivity.

##### Infections

Patients must be monitored closely for infections including tuberculosis before, during and after treatment with Flixabi. Because the elimination of infliximab may take up to six months, monitoring should be continued throughout this period. Further treatment with Flixabi must not be given if a patient develops a serious infection or sepsis.

Caution should be exercised when considering the use of Flixabi in patients with chronic infection or a history of recurrent infections, including concomitant immunosuppressive therapy. Patients should be advised of and avoid exposure to potential risk factors for infection as appropriate.

Tumour necrosis factor alpha (TNF<sub>α</sub>) mediates inflammation and modulates cellular immune responses. Experimental data show that TNF<sub>α</sub> is essential for the clearing of intracellular infections. Clinical experience shows that host defence against infection is compromised in some patients treated with infliximab.

It should be noted that suppression of TNF $\alpha$  may mask symptoms of infection such as fever. Early recognition of atypical clinical presentations of serious infections and of typical clinical presentation of rare and unusual infections is critical in order to minimise delays in diagnosis and treatment.

Patients taking TNF-blockers are more susceptible to serious infections.

Tuberculosis, bacterial infections, including sepsis and pneumonia, invasive fungal, viral, and other opportunistic infections have been observed in patients treated with infliximab. Some of these infections have been fatal; the most frequently reported opportunistic infections with a mortality rate of > 5% include pneumocystosis, candidiasis, listeriosis and aspergillosis.

Patients who develop a new infection while undergoing treatment with Flixabi, should be monitored closely and undergo a complete diagnostic evaluation. Administration of Flixabi should be discontinued if a patient develops a new serious infection or sepsis, and appropriate antimicrobial or antifungal therapy should be initiated until the infection is controlled.

### *Tuberculosis*

There have been reports of active tuberculosis in patients receiving infliximab. It should be noted that in the majority of these reports tuberculosis was extrapulmonary, presenting as either local or disseminated disease.

Before starting treatment with Flixabi, all patients must be evaluated for both active and inactive ('latent') tuberculosis. This evaluation should include a detailed medical history with personal history of tuberculosis or possible previous contact with tuberculosis and previous and/or current immunosuppressive therapy. Appropriate screening tests, i.e. tuberculin skin test and chest X-ray, should be performed in all patients (local recommendations may apply). It is recommended that the conduct of these tests should be recorded in the patient's alert card. Prescribers are reminded of the risk of false negative tuberculin skin test results, especially in patients who are severely ill or immunocompromised.

If active tuberculosis is diagnosed, Flixabi therapy must not be initiated (see section 4.3).

If latent tuberculosis is suspected, a physician with expertise in the treatment of tuberculosis should be consulted. In all situations described below, the benefit/risk balance of Flixabi therapy should be very carefully considered.

If inactive ('latent') tuberculosis is diagnosed, treatment for latent tuberculosis must be started with antituberculosis therapy before the initiation of Flixabi, and in accordance with local recommendations.

In patients who have several or significant risk factors for tuberculosis and have a negative test for latent tuberculosis, antituberculosis therapy should be considered before the initiation of Flixabi.

Use of anti-tuberculosis therapy should also be considered before the initiation of Flixabi in patients with a past history of latent or active tuberculosis in whom an adequate course of treatment cannot be confirmed. Some cases of active tuberculosis have been reported in patients treated with Flixabi during and after treatment for latent tuberculosis.

All patients should be informed to seek medical advice if signs/symptoms suggestive of tuberculosis (e.g. persistent cough, wasting/weight loss, low-grade fever) appear during or after Flixabi treatment.

### *Invasive fungal infections*

In patients treated with Flixabi, an invasive fungal infection such as aspergillosis, candidiasis, pneumocystosis, histoplasmosis, coccidioidomycosis or blastomycosis should be suspected if they develop a serious systemic illness, and a physician with expertise in the diagnosis and treatment of invasive fungal infections should be consulted at an early stage when investigating these patients.

Invasive fungal infections may present as disseminated rather than localised disease, and antigen and antibody testing may be negative in some patients with active infection. Appropriate empiric antifungal therapy should be considered while a diagnostic workup is being performed taking into account both the risk for severe fungal infection and the risks of antifungal therapy.

For patients who have resided in or travelled to regions where invasive fungal infections such as histoplasmosis, coccidioidomycosis, or blastomycosis are endemic, the benefits and risks of Flixabi treatment should be carefully considered before initiation of Flixabi therapy.

#### *Fistulising Crohn's disease*

Patients with fistulising Crohn's disease with acute suppurative fistulas must not initiate Flixabi therapy until a source for possible infection, specifically abscess, has been excluded (see section 4.3).

#### Hepatitis B (HBV) reactivation

Reactivation of hepatitis B has occurred in patients receiving a TNF-antagonist including infliximab, who are chronic carriers of this virus. Some cases have had fatal outcome.

Patients should be tested for HBV infection before initiating treatment with Flixabi. For patients who test positive for HBV infection, consultation with a physician with expertise in the treatment of hepatitis B is recommended. Carriers of HBV who require treatment with Flixabi should be closely monitored for signs and symptoms of active HBV infection throughout therapy and for several months following termination of therapy. Adequate data of treating patients who are carriers of HBV with antiviral therapy in conjunction with TNF-antagonist therapy to prevent HBV reactivation are not available. In patients who develop HBV reactivation, Flixabi should be stopped and effective antiviral therapy with appropriate supportive treatment should be initiated.

#### Hepatobiliary events

Cases of jaundice and non-infectious hepatitis, some with features of autoimmune hepatitis, have been observed in the post-marketing experience of infliximab. Isolated cases of liver failure resulting in liver transplantation or death have occurred. Patients with symptoms or signs of liver dysfunction should be evaluated for evidence of liver injury. If jaundice and/or ALT elevations  $\geq 5$  times the upper limit of normal develop(s), Flixabi should be discontinued, and a thorough investigation of the abnormality should be undertaken.

#### Concurrent administration of TNF-alpha inhibitor and anakinra

Serious infections and neutropenia were seen in clinical studies with concurrent use of anakinra and another TNF $_{\alpha}$ -blocking agent, etanercept, with no added clinical benefit compared to etanercept alone. Because of the nature of the adverse events seen with combination of etanercept and anakinra therapy, similar toxicities may also result from the combination of anakinra and other TNF $_{\alpha}$ -blocking agents. Therefore, the combination of Flixabi and anakinra is not recommended.

#### Concurrent administration of TNF-alpha inhibitor and abatacept

In clinical studies concurrent administration of TNF-antagonists and abatacept has been associated with an increased risk of infections including serious infections compared to TNF-antagonists alone, without increased clinical benefit. The combination of Flixabi and abatacept is not recommended.

#### Concurrent administration with other biological therapeutics

There is insufficient information regarding the concomitant use of infliximab with other biological therapeutics used to treat the same conditions as infliximab. The concomitant use of Flixabi with these biologics is not recommended because of the possibility of an increased risk of infection, and other potential pharmacological interactions.

#### Switching between biological DMARDS

Care should be taken and patients should continue to be monitored when switching from one biologic to another, since overlapping biological activity may further increase the risk for adverse events, including infection.

#### Live vaccines/therapeutic infectious agents

In patients receiving anti-TNF therapy, limited data are available on the response to vaccination with live vaccines or on the secondary transmission of infection by live vaccines. Use of live vaccines can result in

clinical infections, including disseminated infections. The concurrent administration of live vaccines with Flixabi is not recommended.

In infants exposed in utero to infliximab, fatal outcome due to disseminated Bacillus Calmette-Guérin (BCG) infection has been reported following administration of BCG vaccine after birth. At least a six month waiting period following birth is recommended before the administration of live vaccines to infants exposed *in utero* to infliximab (see section 4.6).

Other uses of therapeutic infectious agents such as live attenuated bacteria (e.g., BCG bladder instillation for the treatment of cancer) could result in clinical infections, including disseminated infections. It is recommended that therapeutic infectious agents not be given concurrently with Flixabi.

#### Autoimmune processes

The relative deficiency of TNF $\alpha$  caused by anti-TNF therapy may result in the initiation of an autoimmune process. If a patient develops symptoms suggestive of a lupus-like syndrome following treatment with Flixabi and is positive for antibodies against double-stranded DNA, further treatment with Flixabi must not be given (see section 4.8).

#### Neurological events

Use of TNF-blocking agents, including infliximab, has been associated with cases of new onset or exacerbation of clinical symptoms and/or radiographic evidence of central nervous system demyelinating disorders, including multiple sclerosis, and peripheral demyelinating disorders, including Guillain-Barré syndrome. In patients with pre-existing or recent onset of demyelinating disorders, the benefits and risks of anti-TNF treatment should be carefully considered before initiation of Flixabi therapy. Discontinuation of Flixabi should be considered if these disorders develop.

#### Malignancies and lymphoproliferative disorders

In the controlled portions of clinical studies of TNF-blocking agents, more cases of malignancies including lymphoma have been observed among patients receiving a TNF blocker compared with control patients. During clinical studies of infliximab across all approved indications the incidence of lymphoma in infliximab-treated patients was higher than expected in the general population, but the occurrence of lymphoma was rare. In the post-marketing setting, cases of leukaemia have been reported in patients treated with a TNF-antagonist. There is an increased background risk for lymphoma and leukaemia in rheumatoid arthritis patients with long-standing, highly active, inflammatory disease, which complicates risk estimation.

In an exploratory clinical study evaluating the use of infliximab in patients with moderate to severe chronic obstructive pulmonary disease (COPD), more malignancies were reported in infliximab-treated patients compared with control patients. All patients had a history of heavy smoking. Caution should be exercised in considering treatment of patients with increased risk for malignancy due to heavy smoking.

With the current knowledge, a risk for the development of lymphomas or other malignancies in patients treated with a TNF-blocking agent cannot be excluded (see section 4.8). Caution should be exercised when considering TNF-blocking therapy for patients with a history of malignancy or when considering continuing treatment in patients who develop a malignancy.

Caution should also be exercised in patients with psoriasis and a medical history of extensive immunosuppressant therapy or prolonged PUVA treatment.

Malignancies, some fatal, have been reported among children, adolescents and young adults (up to 22 years of age) treated with TNF-blocking agents (initiation of therapy  $\leq$  18 years of age), including infliximab in the post-marketing setting. Approximately half the cases were lymphomas. The other cases represented a variety of different malignancies and included rare malignancies usually associated with immunosuppression. A risk for the development of malignancies in patients treated with TNF-blockers cannot be excluded.

Post-marketing cases of hepatosplenic T-cell lymphoma (HSTCL) have been reported in patients treated with TNF-blocking agents including infliximab. This rare type of T-cell lymphoma has a very aggressive disease course and is usually fatal. Almost all patients had received treatment with AZA or 6-MP concomitantly with or immediately prior to a TNF-blocker. The vast majority of infliximab cases have occurred in patients with Crohn's disease or ulcerative colitis and most were reported in adolescent or young adult males. The potential risk with the combination of AZA or 6-MP and Flixabi should be carefully considered. A risk for the development for hepatosplenic T-cell lymphoma in patients treated with Flixabi cannot be excluded (see section 4.8).

Melanoma and Merkel cell carcinoma have been reported in patients treated with TNF blocker therapy, including infliximab (see section 4.8). Periodic skin examination is recommended, particularly for patients with risk factors for skin cancer.

A population-based retrospective cohort study using data from Swedish national health registries found an increased incidence of cervical cancer in women with rheumatoid arthritis treated with infliximab compared to biologics-naïve patients or the general population, including those over 60 years of age. Periodic screening should continue in women treated with infliximab, including those over 60 years of age.

All patients with ulcerative colitis who are at increased risk for dysplasia or colon carcinoma (for example, patients with long-standing ulcerative colitis or primary sclerosing cholangitis), or who had a prior history of dysplasia or colon carcinoma should be screened for dysplasia at regular intervals before therapy and throughout their disease course. This evaluation should include colonoscopy and biopsies per local recommendations. With current data it is not known if infliximab treatment influences the risk for developing dysplasia or colon cancer (see section 4.8).

Since the possibility of increased risk of cancer development in patients with newly diagnosed dysplasia treated with infliximab is not established, the risk and benefits to the individual patients must be carefully reviewed and consideration should be given to discontinuation of therapy.

#### Heart failure

Flixabi should be used with caution in patients with mild heart failure (NYHA class I/II). Patients should be closely monitored and Flixabi must not be continued in patients who develop new or worsening symptoms of heart failure (see sections 4.3 and 4.8).

#### Haematologic reactions

There have been reports of pancytopenia, leucopenia, neutropenia, and thrombocytopenia in patients receiving TNF-blockers, including infliximab. All patients should be advised to seek immediate medical attention if they develop signs and symptoms suggestive of blood dyscrasias (e.g. persistent fever, bruising, bleeding, pallor). Discontinuation of Flixabi therapy should be considered in patients with confirmed significant haematologic abnormalities.

#### Others

There is limited safety experience of infliximab treatment in patients who have undergone surgical procedures, including arthroplasty. The long half-life of infliximab should be taken into consideration if a surgical procedure is planned. A patient who requires surgery while on Flixabi should be closely monitored for infections, and appropriate actions should be taken.

Failure to respond to treatment for Crohn's disease may indicate the presence of a fixed fibrotic stricture that may require surgical treatment. There is no evidence to suggest that infliximab worsens or causes fibrotic strictures.

#### Special populations

##### Elderly ( $\geq 65$ years)

The incidence of serious infections in infliximab-treated patients 65 years and older was greater than in those under 65 years of age. Some of those had a fatal outcome. Particular attention regarding the risk for infection should be paid when treating the elderly (see section 4.8).

### Paediatric population

#### *Infections*

In clinical studies, infections have been reported in a higher proportion of paediatric patients compared to adult patients (see section 4.8).

#### *Vaccinations*

It is recommended that paediatric patients, if possible, be brought up to date with all vaccinations in agreement with current vaccination guidelines prior to initiating Flixabi therapy.

#### *Malignancies and lymphoproliferative disorders*

Malignancies, some fatal, have been reported among children, adolescents and young adults (up to 22 years of age) treated with TNF-blocking agents (initiation of therapy  $\leq$  18 years of age), including infliximab in the post-marketing setting. Approximately half the cases were lymphomas. The other cases represented a variety of different malignancies and included rare malignancies usually associated with immunosuppression. A risk for the development of malignancies in children and adolescents treated with TNF-blockers cannot be excluded.

Post-marketing cases of hepatosplenic T-cell lymphoma have been reported in patients treated with TNF-blocking agents including infliximab. This rare type of T-cell lymphoma has a very aggressive disease course and is usually fatal. Almost all patients had received treatment with AZA or 6-MP concomitantly with or immediately prior to a TNF-blocker. The vast majority of infliximab cases have occurred in patients with Crohn's disease or ulcerative colitis and most were reported in adolescent or young adult males. The potential risk with the combination of AZA or 6-MP and Flixabi should be carefully considered. A risk for the development for hepatosplenic T-cell lymphoma in patients treated with Flixabi cannot be excluded (see section 4.8).

## **4.5 Interaction with other medicinal products and other forms of interaction**

No interaction studies have been performed.

In rheumatoid arthritis, psoriatic arthritis and Crohn's disease patients, there are indications that concomitant use of methotrexate and other immunomodulators reduces the formation of antibodies against infliximab and increases the plasma concentrations of infliximab. However, the results are uncertain due to limitations in the methods used for serum analyses of infliximab and antibodies against infliximab.

Corticosteroids do not appear to affect the pharmacokinetics of infliximab to a clinically relevant extent.

The combination of Flixabi with other biological therapeutics used to treat the same conditions as Flixabi, including anakinra and abatacept, is not recommended (see section 4.4).

It is recommended that live vaccines not be given concurrently with Flixabi. It is also recommended that live vaccines not be given to infants after *in utero* exposure to infliximab for at least 6 months following birth (see section 4.4).

It is recommended that therapeutic infectious agents not be given concurrently with Flixabi (see section 4.4).

## **4.6 Fertility, pregnancy and lactation**

### Women of childbearing potential

Women of childbearing potential should consider the use of adequate contraception to prevent pregnancy and continue its use for at least 6 months after the last Flixabi treatment.

### Pregnancy

The moderate number of prospectively collected pregnancies exposed to infliximab resulting in live birth with known outcomes, including approximately 1100 exposed during the first trimester, does not indicate an increase in the rate of malformation in the newborn.

Based on an observational study from Northern Europe, an increased risk (OR, 95% CI; p-value) for C-section (1.50, 1.14-1.96; p=0.0032), preterm birth (1.48, 1.05-2.09; p=0.024), small for gestational age (2.79, 1.54-5.04; p=0.0007), and low birth weight (2.03, 1.41-2.94; p=0.0002) was observed in women exposed during pregnancy to infliximab (with or without immunomodulators/corticosteroids, 270 pregnancies) as compared to women exposed to immunomodulators and/or corticosteroids only (6,460 pregnancies). The potential contribution of exposure to infliximab and/or the severity of the underlying disease in these outcomes remains unclear.

Due to its inhibition of TNF $\alpha$ , infliximab administered during pregnancy could affect normal immune responses in the newborn. In a developmental toxicity study conducted in mice using an analogous antibody that selectively inhibits the functional activity of mouse TNF $\alpha$ , there was no indication of maternal toxicity, embryotoxicity or teratogenicity (see section 5.3).

The available clinical experience is limited. Infliximab should only be used during pregnancy if clearly needed.

Infliximab crosses the placenta and has been detected in the serum of infants up to 6 months following birth. After *in utero* exposure to infliximab, infants may be at increased risk of infection, including serious disseminated infection that can become fatal. Administration of live vaccines (e.g. BCG vaccine) to infants exposed to infliximab *in utero* is not recommended for at least 6 months after birth (see sections 4.4 and 4.5). Cases of agranulocytosis have also been reported (see section 4.8).

### Breast-feeding

It is unknown whether infliximab is excreted in human milk or absorbed systemically after ingestion. Because human immunoglobulins are excreted in milk, women must not breast-feed for at least 6 months after Flixabi treatment.

### Fertility

There are insufficient preclinical data to draw conclusions on the effects of infliximab on fertility and general reproductive function (see section 5.3).

## **4.7 Effects on ability to drive and use machines**

Flixabi may have a minor influence on the ability to drive and use machines. Dizziness may occur following administration of Flixabi (see section 4.8).

## **4.8 Undesirable effects**

### Summary of the safety profile

Upper respiratory tract infection was the most common adverse drug reaction (ADR) reported in clinical trials, occurring in 25.3% of infliximab-treated patients compared with 16.5% of control patients. The most serious ADRs associated with the use of TNF blockers that have been reported for infliximab include HBV reactivation, CHF, serious infections (including sepsis, opportunistic infections and TB), serum sickness (delayed hypersensitivity reactions), haematologic reactions, systemic lupus erythematosus/lupus-like syndrome, demyelinating disorders, hepatobiliary events, lymphoma, HSTCL, leukaemia, Merkel cell carcinoma, melanoma, paediatric malignancy, sarcoidosis/sarcoid-like reaction, intestinal or perianal abscess (in Crohn's disease), and serious infusion reactions (see section 4.4).

Tabulated list of adverse reactions

Table 1 lists ADRs based on experience from clinical studies as well as adverse reactions, some with fatal outcome, reported from post-marketing experience. Within the organ system classes, adverse reactions are listed under headings of frequency using the following categories: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ), not known (cannot be estimated from the available data). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

**Table 1**  
**Undesirable effects in clinical studies and from post-marketing experience**

Infections and infestations	
Very Common:	Viral infection (e.g. influenza, herpes virus infection).
Common:	Bacterial infections (e.g. sepsis, cellulitis, abscess).
Uncommon:	Tuberculosis, fungal infections (e.g. candidiasis).
Rare:	Meningitis, opportunistic infections (such as invasive fungal infections [pneumocystosis, histoplasmosis, aspergillosis, coccidioidomycosis, cryptococcosis, blastomycosis], bacterial infections [atypical mycobacterial, listeriosis, salmonellosis], and viral infections [cytomegalovirus]), parasitic infections, hepatitis B reactivation.
Not known:	Vaccine breakthrough infection (after <i>in utero</i> exposure to infliximab)*.
Neoplasms benign, malignant and unspecified (including cysts and polyps)	
Rare:	Lymphoma, non-Hodgkin's lymphoma, Hodgkin's disease, leukaemia, melanoma, cervical cancer.
Not known:	Hepatosplenic T-cell lymphoma (primarily in adolescents and young adults with Crohn's disease and ulcerative colitis), Merkel cell carcinoma.
Blood and lymphatic system disorders	
Common:	Neutropenia, leucopenia, anaemia, lymphadenopathy.
Uncommon:	Thrombocytopenia, lymphopenia, lymphocytosis.
Rare:	Agranulocytosis (including infants exposed <i>in utero</i> to infliximab), thrombotic thrombocytopenic purpura, pancytopenia, haemolytic anaemia, idiopathic thrombocytopenic purpura.
Immune system disorders	
Common:	Allergic respiratory symptom.
Uncommon:	Anaphylactic reaction, lupus-like syndrome, serum sickness or serum sickness-like reaction.
Rare:	Anaphylactic shock, vasculitis, sarcoid-like reaction.
Psychiatric disorders	
Common:	Depression, insomnia.
Uncommon:	Amnesia, agitation, confusion, somnolence, nervousness.
Rare:	Apathy.
Nervous system disorders	
Very common:	Headache.
Common:	Vertigo, dizziness, hypoaesthesia, paraesthesia.
Uncommon:	Seizure, neuropathy.
Rare:	Transverse myelitis, central nervous system demyelinating disorders (multiple sclerosis-like disease and optic neuritis), peripheral demyelinating disorders (such as Guillain-Barré syndrome, chronic inflammatory demyelinating polyneuropathy and multifocal motor neuropathy).

Eye disorders	Common:	Conjunctivitis.
	Uncommon:	Keratitis, periorbital oedema, hordeolum.
	Rare:	Endophthalmitis.
	Not known:	Transient visual loss occurring during or within 2 hours of infusion.
Cardiac disorders	Common:	Tachycardia, palpitation.
	Uncommon:	Cardiac failure (new onset or worsening), arrhythmia, syncope, bradycardia.
	Rare:	Cyanosis, pericardial effusion.
	Not known:	Myocardial ischaemia/myocardial infarction.
Vascular disorders	Common:	Hypotension, hypertension, ecchymosis, hot flush, flushing.
	Uncommon:	Peripheral ischaemia, thrombophlebitis, haematoma.
	Rare:	Circulatory failure, petechia, vasospasm.
Respiratory, thoracic and mediastinal disorders	Very common:	Upper respiratory tract infection, sinusitis.
	Common:	Lower respiratory tract infection (e.g. bronchitis, pneumonia), dyspnoea, epistaxis.
	Uncommon:	Pulmonary oedema, bronchospasm, pleurisy, pleural effusion.
	Rare:	Interstitial lung disease (including rapidly progressive disease, lung fibrosis and pneumonitis).
Gastrointestinal disorders	Very common:	Abdominal pain, nausea.
	Common:	Gastrointestinal haemorrhage, diarrhoea, dyspepsia, gastroesophageal reflux, constipation.
	Uncommon:	Intestinal perforation, intestinal stenosis, diverticulitis, pancreatitis, cheilitis.
Hepatobiliary disorders	Common:	Hepatic function abnormal, transaminases increased.
	Uncommon:	Hepatitis, hepatocellular damage, cholecystitis.
	Rare:	Autoimmune hepatitis, jaundice.
	Not known:	Liver failure.
Skin and subcutaneous tissue disorders	Common:	New onset or worsening psoriasis including pustular psoriasis (primarily palm & soles), urticaria, rash, pruritus, hyperhidrosis, dry skin, fungal dermatitis, eczema, alopecia.
	Uncommon:	Bullous eruption, onychomycosis, seborrhoea, rosacea, skin papilloma, hyperkeratosis, abnormal skin pigmentation.
	Rare:	Toxic epidermal necrolysis, Stevens-Johnson Syndrome, erythema multiforme, furunculosis.
	Not known:	Worsening of symptoms of dermatomyositis.
Musculoskeletal and connective tissue disorders	Common:	Arthralgia, myalgia, back pain.
Renal and urinary disorders	Common:	Urinary tract infection.
	Uncommon:	Pyelonephritis.
Reproductive system and breast disorders	Uncommon:	Vaginitis.
General disorders and administration site conditions	Very common:	Infusion-related reaction, pain.
	Common:	Chest pain, fatigue, fever, injection site reaction, chills, oedema.
	Uncommon:	Impaired healing.
	Rare:	Granulomatous lesion.
Investigations		

Uncommon:	Autoantibody positive.
Rare:	Complement factor abnormal.

\* including bovine tuberculosis (disseminated BCG infection), see section 4.4

## Description of selected adverse reactions

### Infusion-related reactions

An infusion-related reaction was defined in clinical studies as any adverse event occurring during an infusion or within 1 hour after an infusion. In Phase III clinical studies, 18% of infliximab-treated patients compared with 5% of placebo-treated patients experienced an infusion-related reaction. Overall, a higher proportion of patients receiving infliximab monotherapy experienced an infusion-related reaction compared to patients receiving infliximab with concomitant immunomodulators. Approximately 3% of patients discontinued treatment due to infusion-related reactions and all patients recovered with or without medical therapy. Of infliximab-treated patients who had an infusion reaction during the induction period, through week 6, 27% experienced an infusion reaction during the maintenance period, week 7 through week 54. Of patients who did not have an infusion reaction during the induction period, 9% experienced an infusion reaction during the maintenance period.

In a clinical study of patients with rheumatoid arthritis (ASPIRE), infusions were to be administered over 2 hours for the first 3 infusions. The duration of subsequent infusions could be shortened to not less than 40 minutes in patients who did not experience serious infusion reactions. In this trial, sixty six percent of the patients (686 out of 1,040) received at least one shortened infusion of 90 minutes or less and 44% of the patients (454 out of 1,040) received at least one shortened infusion of 60 minutes or less. Of the infliximab-treated patients who received at least one shortened infusion, infusion-related reactions occurred in 15% of patients and serious infusion reactions occurred in 0.4% of patients.

In a clinical study of patients with Crohn's disease (SONIC), infusion-related reactions occurred in 16.6% (27/163) of patients receiving infliximab monotherapy, 5% (9/179) of patients receiving infliximab in combination with AZA, and 5.6% (9/161) of patients receiving AZA monotherapy. One serious infusion reaction (< 1%) occurred in a patient on infliximab monotherapy.

In post-marketing experience, cases of anaphylactic-like reactions, including laryngeal/pharyngeal oedema and severe bronchospasm, and seizure have been associated with infliximab administration (see section 4.4). Cases of transient visual loss occurring during or within 2 hours of infliximab infusion have been reported. Events (some fatal) of myocardial ischaemia/infarction and arrhythmia have also been reported, some in close temporal association with infusion of infliximab.

### Infusion reactions following re-administration of infliximab

A clinical study in patients with moderate to severe psoriasis was designed to assess the efficacy and safety of long-term maintenance therapy versus re-treatment with an induction regimen of infliximab (maximum of four infusions at 0, 2, 6 and 14 weeks) following disease flare. Patients did not receive any concomitant immunosuppressant therapy. In the re-treatment arm, 4% (8/219) of patients experienced a serious infusion reaction versus < 1% (1/222) on maintenance therapy. The majority of serious infusion reactions occurred during the second infusion at week 2. The interval between the last maintenance dose and the first re-induction dose ranged from 35-231 days. Symptoms included, but were not limited to, dyspnoea, urticaria, facial oedema, and hypotension. In all cases, infliximab treatment was discontinued and/or other treatment instituted with complete resolution of signs and symptoms.

### Delayed hypersensitivity

In clinical studies delayed hypersensitivity reactions have been uncommon and have occurred after infliximab-free intervals of less than 1 year. In the psoriasis studies, delayed hypersensitivity reactions occurred early in the treatment course. Signs and symptoms included myalgia and/or arthralgia with fever and/or rash, with some patients experiencing pruritus, facial, hand or lip oedema, dysphagia, urticaria, sore throat and headache.

There are insufficient data on the incidence of delayed hypersensitivity reactions after infliximab-free intervals of more than 1 year but limited data from clinical studies suggest an increased risk for delayed hypersensitivity with increasing infliximab-free interval (see section 4.4).

In a 1-year clinical study with repeated infusions in patients with Crohn's disease (ACCENT I study), the incidence of serum sickness-like reactions was 2.4%.

#### Immunogenicity

Patients who developed antibodies to infliximab were more likely (approximately 2-3 fold) to develop infusion-related reactions. Use of concomitant immunosuppressant agents appeared to reduce the frequency of infusion-related reactions.

In clinical studies using single and multiple infliximab doses ranging from 1 to 20 mg/kg, antibodies to infliximab were detected in 14% of patients with any immunosuppressant therapy, and in 24% of patients without immunosuppressant therapy. In rheumatoid arthritis patients who received the recommended repeated treatment dose regimens with methotrexate, 8% of patients developed antibodies to infliximab. In psoriatic arthritis patients who received 5 mg/kg with and without methotrexate, antibodies occurred overall in 15% of patients (antibodies occurred in 4% of patients receiving methotrexate and in 26% of patients not receiving methotrexate at baseline). In Crohn's disease patients who received maintenance treatment, antibodies to infliximab occurred overall in 3.3% of patients receiving immunosuppressants and in 13.3% of patients not receiving immunosuppressants. The antibody incidence was 2-3 fold higher for patients treated episodically. Due to methodological limitations, a negative assay did not exclude the presence of antibodies to infliximab. Some patients who developed high titres of antibodies to infliximab had evidence of reduced efficacy. In psoriasis patients treated with infliximab as a maintenance regimen in the absence of concomitant immunomodulators, approximately 28% developed antibodies to infliximab (see section 4.4: "Infusion reactions and hypersensitivity").

#### Infections

Tuberculosis, bacterial infections, including sepsis and pneumonia, invasive fungal, viral, and other opportunistic infections have been observed in patients receiving infliximab. Some of these infections have been fatal; the most frequently reported opportunistic infections with a mortality rate of > 5% include pneumocystosis, candidiasis, listeriosis and aspergillosis (see section 4.4).

In clinical studies 36% of infliximab-treated patients were treated for infections compared with 25% of placebo-treated patients.

In rheumatoid arthritis clinical studies, the incidence of serious infections including pneumonia was higher in infliximab plus methotrexate-treated patients compared with methotrexate alone especially at doses of 6 mg/kg or greater (see section 4.4).

In post-marketing spontaneous reporting, infections are the most common serious adverse event. Some of the cases have resulted in a fatal outcome. Nearly 50% of reported deaths have been associated with infection. Cases of tuberculosis, sometimes fatal, including miliary tuberculosis and tuberculosis with extra-pulmonary location have been reported (see section 4.4).

#### Malignancies and lymphoproliferative disorders

In clinical studies with infliximab in which 5,780 patients were treated, representing 5,494 patient years, 5 cases of lymphomas and 26 non-lymphoma malignancies were detected as compared with no lymphomas and 1 non-lymphoma malignancy in 1,600 placebo-treated patients representing 941 patient years.

In long-term safety follow-up of clinical studies with infliximab of up to 5 years, representing 6,234 patients-years (3,210 patients), 5 cases of lymphoma and 38 cases of non-lymphoma malignancies were reported.

Cases of malignancies, including lymphoma, have also been reported in the post-marketing setting (see section 4.4).

In an exploratory clinical study involving patients with moderate to severe COPD who were either current smokers or ex-smokers, 157 adult patients were treated with infliximab at doses similar to those used in rheumatoid arthritis and Crohn’s disease. Nine of these patients developed malignancies, including 1 lymphoma. The median duration of follow-up was 0.8 years (incidence 5.7% [95% CI 2.65%-10.6%]). There was one reported malignancy amongst 77 control patients (median duration of follow-up 0.8 years; incidence 1.3% [95% CI 0.03%-7.0%]). The majority of the malignancies developed in the lung or head and neck.

A population-based retrospective cohort study found an increased incidence of cervical cancer in women with rheumatoid arthritis treated with infliximab compared to biologics-naïve patients or the general population, including those over 60 years of age (see section 4.4).

In addition, post-marketing cases of hepatosplenic T-cell lymphoma have been reported in patients treated with infliximab with the vast majority of cases occurring in Crohn’s disease and ulcerative colitis, and most of whom were adolescent or young adult males (see section 4.4).

#### Heart failure

In a Phase II study aimed at evaluating infliximab in congestive heart failure (CHF), higher incidence of mortality due to worsening of heart failure were seen in patients treated with infliximab, especially those treated with the higher dose of 10 mg/kg (i.e. twice the maximum approved dose). In this study 150 patients with NYHA Class III-IV CHF (left ventricular ejection fraction  $\leq$  35%) were treated with 3 infusions of infliximab 5 mg/kg, 10 mg/kg, or placebo over 6 weeks. At 38 weeks, 9 of 101 patients treated with infliximab (2 at 5 mg/kg and 7 at 10 mg/kg) died compared to one death among the 49 patients on placebo. There have been post-marketing reports of worsening heart failure, with and without identifiable precipitating factors, in patients taking infliximab. There have also been post-marketing reports of new onset heart failure, including heart failure in patients without known pre-existing cardiovascular disease. Some of these patients have been under 50 years of age.

#### Hepatobiliary events

In clinical studies, mild or moderate elevations of ALT and AST have been observed in patients receiving infliximab without progression to severe hepatic injury. Elevations of ALT  $\geq$  5 x Upper Limit of Normal (ULN) have been observed (see Table 2). Elevations of aminotransferases were observed (ALT more common than AST) in a greater proportion of patients receiving infliximab than in controls, both when infliximab was given as monotherapy and when it was used in combination with other immunosuppressive agents. Most aminotransferase abnormalities were transient; however, a small number of patients experienced more prolonged elevations. In general, patients who developed ALT and AST elevations were asymptomatic, and the abnormalities decreased or resolved with either continuation or discontinuation of infliximab, or modification of concomitant therapy. In post-marketing surveillance, cases of jaundice and hepatitis, some with features of autoimmune hepatitis, have been reported in patients receiving infliximab (see section 4.4).

**Table 2**  
**Proportion of patients with increased ALT activity in clinical studies**

Indication	Number of patients <sup>3</sup>		Median follow-up (wks) <sup>4</sup>		$\geq$ 3 x ULN		$\geq$ 5 x ULN	
	placebo	infliximab	placebo	infliximab	placebo	infliximab	placebo	infliximab
Rheumatoid arthritis <sup>1</sup>	375	1,087	58.1	58.3	3.2%	3.9%	0.8%	0.9%
Crohn’s disease <sup>2</sup>	324	1034	53.7	54.0	2.2%	4.9%	0.0%	1.5%
Paediatric Crohn’s disease	N/A	139	N/A	53.0	N/A	4.4%	N/A	1.5%
Ulcerative colitis	242	482	30.1	30.8	1.2%	2.5%	0.4%	0.6%

Indication	Number of patients <sup>3</sup>		Median follow-up (wks) <sup>4</sup>		≥ 3 x ULN		≥ 5 x ULN	
	placebo	infliximab	placebo	infliximab	placebo	infliximab	placebo	infliximab
Paediatric Ulcerative colitis	N/A	60	N/A	49.4	N/A	6.7%	N/A	1.7%
Ankylosing spondylitis	76	275	24.1	101.9	0.0%	9.5%	0.0%	3.6%
Psoriatic arthritis	98	191	18.1	39.1	0.0%	6.8%	0.0%	2.1%
Plaque psoriasis	281	1,175	16.1	50.1	0.4%	7.7%	0.0%	3.4%

<sup>1</sup> Placebo patients received methotrexate while infliximab patients received both infliximab and methotrexate.

<sup>2</sup> Placebo patients in the 2 Phase III studies in Crohn's disease, ACCENT I and ACCENT II, received an initial dose of 5 mg/kg infliximab at study start and were on placebo in the maintenance phase. Patients who were randomised to the placebo maintenance group and then later crossed over to infliximab are included in the infliximab group in the ALT analysis. In the Phase IIIb trial in Crohn's disease, SONIC, placebo patients received AZA 2.5 mg/kg/day as active control in addition to placebo infliximab infusions.

<sup>3</sup> Number of patients evaluated for ALT.

<sup>4</sup> Median follow-up is based on patients treated.

### Antinuclear antibodies (ANA)/Anti-double-stranded DNA (dsDNA) antibodies

Approximately half of infliximab-treated patients in clinical studies who were ANA negative at baseline developed a positive ANA during the study compared with approximately one fifth of placebo-treated patients. Anti-dsDNA antibodies were newly detected in approximately 17% of infliximab-treated patients compared with 0% of placebo-treated patients. At the last evaluation, 57% of infliximab-treated patients remained anti-dsDNA positive. Reports of lupus and lupus-like syndromes, however, remain uncommon (see section 4.4).

### Paediatric population

#### Juvenile rheumatoid arthritis patients

Infliximab was studied in a clinical study in 120 patients (age range: 4-17 years old) with active juvenile rheumatoid arthritis despite methotrexate. Patients received 3 or 6 mg/kg infliximab as a 3-dose induction regimen (weeks 0, 2, 6 or weeks 14, 16, 20, respectively) followed by maintenance therapy every 8 weeks, in combination with methotrexate.

#### Infusion reactions

Infusion reactions occurred in 35% of patients with juvenile rheumatoid arthritis receiving 3 mg/kg compared with 17.5% of patients receiving 6 mg/kg. In the 3 mg/kg infliximab group, 4 out of 60 patients had a serious infusion reaction and 3 patients reported a possible anaphylactic reaction (2 of which were among the serious infusion reactions). In the 6 mg/kg group, 2 out of 57 patients had a serious infusion reaction, one of whom had a possible anaphylactic reaction (see section 4.4).

#### Immunogenicity

Antibodies to infliximab developed in 38% of patients receiving 3 mg/kg compared with 12% of patients receiving 6 mg/kg. The antibody titres were notably higher for the 3 mg/kg compared to the 6 mg/kg group.

#### Infections

Infections occurred in 68% (41/60) of children receiving 3 mg/kg over 52 weeks, 65% (37/57) of children receiving infliximab 6 mg/kg over 38 weeks and 47% (28/60) of children receiving placebo over 14 weeks (see section 4.4).

#### Paediatric Crohn's disease patients

The following adverse events were reported more commonly in paediatric Crohn's disease patients in the REACH study (see section 5.1) than in adult Crohn's disease patients: anaemia (10.7%), blood in stool (9.7%), leucopenia (8.7%), flushing (8.7%), viral infection (7.8%), neutropenia (6.8%), bone fracture (6.8%),

bacterial infection (5.8%), and respiratory tract allergic reaction (5.8%). Other special considerations are discussed below.

#### *Infusion-related reactions*

In REACH, 17.5% of randomised patients experienced 1 or more infusion reactions. There were no serious infusion reactions, and 2 subjects in REACH had non-serious anaphylactic reactions.

#### *Immunogenicity*

Antibodies to infliximab were detected in 3 (2.9%) paediatric patients.

#### *Infections*

In the REACH study, infections were reported in 56.3% of randomised subjects treated with infliximab. Infections were reported more frequently for subjects who received q8 week as opposed to q12 week infusions (73.6% and 38.0%, respectively), while serious infections were reported for 3 subjects in the q8 week and 4 subjects in the q12 week maintenance treatment group. The most commonly reported infections were upper respiratory tract infection and pharyngitis, and the most commonly reported serious infection was abscess. Three cases of pneumonia (1 serious) and 2 cases of herpes zoster (both non-serious) were reported.

#### *Paediatric ulcerative colitis patients*

Overall, the adverse reactions reported in the paediatric ulcerative colitis trial (C0168T72) and adult ulcerative colitis (ACT 1 and ACT 2) studies were generally consistent. In C0168T72, the most common adverse reactions were upper respiratory tract infection, pharyngitis, abdominal pain, fever, and headache. The most common adverse event was worsening of ulcerative colitis, the incidence of which was higher in patients on the q12 week vs. the q8 week dosing regimen.

#### *Infusion-related reactions*

Overall, 8 (13.3%) of 60 treated patients experienced one or more infusion reactions, with 4 of 22 (18.2%) in the q8 week and 3 of 23 (13.0%) in the q12 week treatment maintenance group. No serious infusion reactions were reported. All infusion reactions were mild or moderate in intensity.

#### *Immunogenicity*

Antibodies to infliximab were detected in 4 (7.7%) patients through week 54.

#### *Infections*

Infections were reported in 31 (51.7%) of 60 treated patients in C0168T72 and 22 (36.7%) required oral or parenteral antimicrobial treatment. The proportion of patients with infections in C0168T72 was similar to that in the paediatric Crohn's disease study (REACH) but higher than the proportion in the adults ulcerative colitis studies (ACT 1 and ACT 2). The overall incidence of infections in C0168T72 was 13/22 (59%) in the every 8 week maintenance treatment group and 14/23 (60.9%) in the every 12 week maintenance treatment group. Upper respiratory tract infection (7/60 [12%]) and pharyngitis (5/60 [8%]) were the most frequently reported respiratory system infections. Serious infections were reported in 12% (7/60) of all treated patients.

In this study, there were more patients in the 12 to 17 year age group than in the 6 to 11 year age group (45/60 [75.0%]) vs. 15/60 [25.0%]). While the numbers of patients in each subgroup are too small to make any definitive conclusions about the effect of age on safety events, there were higher proportions of patients with serious adverse events and discontinuation due to adverse events in the younger age group than in the older age group. While the proportion of patients with infections was also higher in the younger age group, for serious infections, the proportions were similar in the two age groups. Overall proportions of adverse events and infusion reactions were similar between the 6 to 11 and 12 to 17 year age groups.

#### Post-marketing experience

Post-marketing spontaneous serious adverse events with infliximab in the paediatric population have included malignancies including hepatosplenic T-cell lymphomas, transient hepatic enzyme abnormalities, lupus-like syndromes, and positive auto-antibodies (see sections 4.4 and 4.8).

#### Additional information on special populations

##### *Elderly (≥ 65 years)*

In rheumatoid arthritis clinical studies, the incidence of serious infections was greater in infliximab plus methotrexate-treated patients 65 years and older (11.3%) than in those under 65 years of age (4.6%). In patients treated with methotrexate alone, the incidence of serious infections was 5.2% in patients 65 years and older compared to 2.7% in patients under 65 (see section 4.4).

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

## **4.9 Overdose**

No case of overdose has been reported. Single doses up to 20 mg/kg have been administered without toxic effects.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: immunosuppressants, tumour necrosis factor alpha (TNF<sub>α</sub>) inhibitors, ATC code: L04AB02.

Flixabi is a biosimilar medicinal product. Detailed information is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

#### Mechanism of action

Infliximab is a chimeric human-murine monoclonal antibody that binds with high affinity to both soluble and transmembrane forms of TNF<sub>α</sub> but not to lymphotoxin α (TNF<sub>β</sub>).

#### Pharmacodynamic effects

Infliximab inhibits the functional activity of TNF<sub>α</sub> in a wide variety of *in vitro* bioassays. Infliximab prevented disease in transgenic mice that develop polyarthritis as a result of constitutive expression of human TNF<sub>α</sub> and when administered after disease onset, it allowed eroded joints to heal. *In vivo*, infliximab rapidly forms stable complexes with human TNF<sub>α</sub>, a process that parallels the loss of TNF<sub>α</sub> bioactivity.

Elevated concentrations of TNF<sub>α</sub> have been found in the joints of rheumatoid arthritis patients and correlate with elevated disease activity. In rheumatoid arthritis, treatment with infliximab reduced infiltration of inflammatory cells into inflamed areas of the joint as well as expression of molecules mediating cellular adhesion, chemoattraction and tissue degradation. After infliximab treatment, patients exhibited decreased levels of serum interleukin 6 (IL-6) and C-reactive protein (CRP), and increased haemoglobin levels in rheumatoid arthritis patients with reduced haemoglobin levels, compared with baseline. Peripheral blood lymphocytes further showed no significant decrease in number or in proliferative responses to *in vitro* mitogenic stimulation when compared with untreated patients' cells. In psoriasis patients, treatment with infliximab resulted in decreases in epidermal inflammation and normalisation of keratinocyte differentiation in psoriatic plaques. In psoriatic arthritis, short term treatment with infliximab reduced the number of T-cells and blood vessels in the synovium and psoriatic skin.

Histological evaluation of colonic biopsies, obtained before and 4 weeks after administration of infliximab, revealed a substantial reduction in detectable TNF<sub>α</sub>. Infliximab treatment of Crohn's disease patients was

also associated with a substantial reduction of the commonly elevated serum inflammatory marker, CRP. Total peripheral white blood cell counts were minimally affected in infliximab-treated patients, although changes in lymphocytes, monocytes and neutrophils reflected shifts towards normal ranges. Peripheral blood mononuclear cells (PBMC) from infliximab-treated patients showed undiminished proliferative responsiveness to stimuli compared with untreated patients, and no substantial changes in cytokine production by stimulated PBMC were observed following treatment with infliximab. Analysis of lamina propria mononuclear cells obtained by biopsy of the intestinal mucosa showed that infliximab treatment caused a reduction in the number of cells capable of expressing TNF $\alpha$  and interferon  $\gamma$ . Additional histological studies provided evidence that treatment with infliximab reduces the infiltration of inflammatory cells into affected areas of the intestine and the presence of inflammation markers at these sites. Endoscopic studies of intestinal mucosa have shown evidence of mucosal healing in infliximab-treated patients.

### Clinical efficacy and safety

#### Adult rheumatoid arthritis

The efficacy of infliximab was assessed in two multicentre, randomised, double-blind, pivotal clinical studies: ATTRACT and ASPIRE. In both studies concurrent use of stable doses of folic acid, oral corticosteroids ( $\leq 10$  mg/day) and/or non-steroidal anti-inflammatory drugs (NSAIDs) was permitted.

The primary endpoints were the reduction of signs and symptoms as assessed by the American College of Rheumatology criteria (ACR20 for ATTRACT, landmark ACR-N for ASPIRE), the prevention of structural joint damage, and the improvement in physical function. A reduction in signs and symptoms was defined to be at least a 20% improvement (ACR20) in both tender and swollen joint counts, and in 3 of the following 5 criteria: (1) evaluator's global assessment, (2) patient's global assessment, (3) functional/disability measure, (4) visual analogue pain scale and (5) erythrocyte sedimentation rate or C-reactive protein. ACR-N uses the same criteria as the ACR20, calculated by taking the lowest percent improvement in swollen joint count, tender joint count, and the median of the remaining 5 components of the ACR response. Structural joint damage (erosions and joint space narrowing) in both hands and feet was measured by the change from baseline in the total van der Heijde-modified Sharp score (0-440). The Health Assessment Questionnaire (HAQ; scale 0-3) was used to measure patients' average change from baseline scores over time, in physical function.

The ATTRACT study evaluated responses at 30, 54 and 102 weeks in a placebo-controlled study of 428 patients with active rheumatoid arthritis despite treatment with methotrexate. Approximately 50% of patients were in functional Class III. Patients received placebo, 3 mg/kg or 10 mg/kg infliximab at weeks 0, 2 and 6, and then every 4 or 8 weeks thereafter. All patients were on stable methotrexate doses (median 15 mg/wk) for 6 months prior to enrolment and were to remain on stable doses throughout the study. Results from week 54 (ACR20, total van der Heijde-modified Sharp score and HAQ) are shown in Table 3. Higher degrees of clinical response (ACR50 and ACR70) were observed in all infliximab groups at 30 and 54 weeks compared with methotrexate alone.

A reduction in the rate of the progression of structural joint damage (erosions and joint space narrowing) was observed in all infliximab groups at 54 weeks (Table 3).

The effects observed at 54 weeks were maintained through 102 weeks. Due to a number of treatment withdrawals, the magnitude of the effect difference between infliximab and the methotrexate alone group cannot be defined.

**Table 3**  
**Effects on ACR20, Structural Joint Damage and Physical Function at week 54, ATTRACT**

	Control <sup>a</sup>	Infliximab <sup>b</sup>				
		3 mg/kg q 8 wks	3 mg/kg q 4 wks	10 mg/kg q 8 wks	10 mg/kg q 4 wks	All infliximab <sup>b</sup>
Patients with ACR20 response/ Patients evaluated (%)	15/88 (17%)	36/86 (42%)	41/86 (48%)	51/87 (59%)	48/81 (59%)	176/340 (52%)

	Infliximab <sup>b</sup>					
	Control <sup>a</sup>	3 mg/kg q 8 wks	3 mg/kg q 4 wks	10 mg/kg q 8 wks	10 mg/kg q 4 wks	All infliximab <sup>b</sup>
Total score <sup>d</sup> (van der Heijde-modified Sharp score)						
Change from baseline (Mean ± SD <sup>c</sup> )	7.0 ± 10.3	1.3 ± 6.0	1.6 ± 8.5	0.2 ± 3.6	-0.7 ± 3.8	0.6 ± 5.9
Median (Interquartile range)	4.0 (0.5,9.7)	0.5 (-1.5,3.0)	0.1 (-2.5,3.0)	0.5 (-1.5,2.0)	-0.5 (-3.0,1.5)	0.0 (-1.8,2.0)
Patients with no deterioration/patients evaluated (%) <sup>c</sup>	13/64 (20%)	34/71 (48%)	35/71 (49%)	37/77 (48%)	44/66 (67%)	150/285 (53%)
HAQ change from baseline over time <sup>e</sup> (patients evaluated)	87	86	85	87	81	339
Mean ± SD <sup>c</sup>	0.2 ± 0.3	0.4 ± 0.3	0.5 ± 0.4	0.5 ± 0.5	0.4 ± 0.4	0.4 ± 0.4

<sup>a</sup> control = All patients had active RA despite treatment with stable methotrexate doses for 6 months prior to enrolment and were to remain on stable doses throughout the study. Concurrent use of stable doses of oral corticosteroids ( $\leq 10$  mg/day) and/or NSAIDs was permitted, and folate supplementation was given.

<sup>b</sup> all infliximab doses given in combination with methotrexate and folate with some on corticosteroids and/or NSAIDs

<sup>c</sup>  $p < 0.001$ , for each infliximab treatment group vs. control

<sup>d</sup> greater values indicate more joint damage.

<sup>e</sup> HAQ = Health Assessment Questionnaire; greater values indicate less disability.

The ASPIRE study evaluated responses at 54 weeks in 1,004 methotrexate naive patients with early ( $\leq 3$  years disease duration, median 0.6 years) active rheumatoid arthritis (median swollen and tender joint count of 19 and 31, respectively). All patients received methotrexate (optimised to 20 mg/wk by week 8) and either placebo, 3 mg/kg or 6 mg/kg infliximab at weeks 0, 2, and 6 and every 8 weeks thereafter. Results from week 54 are shown in Table 4.

After 54 weeks of treatment, both doses of infliximab + methotrexate resulted in statistically significantly greater improvement in signs and symptoms compared to methotrexate alone as measured by the proportion of patients achieving ACR20, 50 and 70 responses.

In ASPIRE, more than 90% of patients had at least two evaluable X-rays. Reduction in the rate of progression of structural damage was observed at weeks 30 and 54 in the infliximab + methotrexate groups compared to methotrexate alone.

**Table 4**  
**Effects on ACRn, Structural Joint Damage and Physical Function at week 54, ASPIRE**

	Infliximab + MTX			
	Placebo + MTX	3 mg/kg	6 mg/kg	Combined
Subjects randomised	282	359	363	722
Percentage ACR improvement				
Mean ± SD <sup>a</sup>	24.8 ± 59.7	37.3 ± 52.8	42.0 ± 47.3	39.6 ± 50.1
Change from baseline in total van der Heijde-modified Sharp score <sup>b</sup>				
Mean ± SD <sup>a</sup>	3.70 ± 9.61	0.42 ± 5.82	0.51 ± 5.55	0.46 ± 5.68
Median	0.43	0.00	0.00	0.00
Improvement from baseline in HAQ averaged over time from week 30 to week 54				
Mean ± SD <sup>d</sup>	0.68 ± 0.63	0.80 ± 0.65	0.88 ± 0.65	0.84 ± 0.65

<sup>a</sup>  $p < 0.001$ , for each infliximab treatment group vs control

<sup>b</sup> greater values indicate more joint damage.

<sup>c</sup> HAQ = Health Assessment Questionnaire; greater values indicate less disability.

<sup>d</sup>  $p = 0.030$  and  $< 0.001$  for the 3 mg/kg and 6 mg/kg treatment groups respectively vs. placebo + MTX.

Data to support dose titration in rheumatoid arthritis come from ATTRACT, ASPIRE and the START study. START was a randomised, multicentre, double-blind, 3-arm, parallel-group safety study. In one of the study arms (group 2, n=329), patients with an inadequate response were allowed to dose titrate with 1.5 mg/kg increments from 3 up to 9 mg/kg. The majority (67%) of these patients did not require any dose titration. Of the patients who required a dose titration, 80% achieved clinical response and the majority (64%) of these required only one adjustment of 1.5 mg/kg.

#### Adult Crohn's disease

##### *Induction treatment in moderately to severely active Crohn's disease*

The efficacy of a single dose treatment with infliximab was assessed in 108 patients with active Crohn's disease (Crohn's Disease Activity Index (CDAI)  $\geq 220 \leq 400$ ) in a randomised, double-blinded, placebo-controlled, dose-response study. Of these 108 patients, 27 were treated with the recommended dosage of infliximab 5 mg/kg. All patients had experienced an inadequate response to prior conventional therapies. Concurrent use of stable doses of conventional therapies was permitted, and 92% of patients continued to receive these therapies.

The primary endpoint was the proportion of patients who experienced a clinical response, defined as a decrease in CDAI by  $\geq 70$  points from baseline at the 4-week evaluation and without an increase in the use of medicinal products or surgery for Crohn's disease. Patients who responded at week 4 were followed to week 12. Secondary endpoints included the proportion of patients in clinical remission at week 4 (CDAI  $< 150$ ) and clinical response over time.

At week 4, following administration of a single dose, 22/27 (81%) of infliximab-treated patients receiving a 5 mg/kg dose achieved a clinical response vs. 4/25 (16%) of the placebo-treated patients ( $p < 0.001$ ). Also at week 4, 13/27 (48%) of infliximab-treated patients achieved a clinical remission (CDAI  $< 150$ ) vs. 1/25 (4%) of placebo-treated patients. A response was observed within 2 weeks, with a maximum response at 4 weeks. At the last observation at 12 weeks, 13/27 (48%) of infliximab-treated patients were still responding.

##### *Maintenance treatment in moderately to severely active Crohn's disease in adults*

The efficacy of repeated infusions with infliximab was studied in a 1-year clinical study (ACCENT I). A total of 573 patients with moderately to severely active Crohn's disease (CDAI  $\geq 220 \leq 400$ ) received a single infusion of 5 mg/kg at week 0. 178 of the 580 enrolled patients (30.7%) were defined as having severe disease (CDAI score  $> 300$  and concomitant corticosteroid and/or immunosuppressants) corresponding to the population defined in the indication (see section 4.1). At week 2, all patients were assessed for clinical response and randomised to one of 3 treatment groups; a placebo maintenance group, 5 mg/kg maintenance group and 10 mg/kg maintenance group. All 3 groups received repeated infusions at week 2, 6 and every 8 weeks thereafter.

Of the 573 patients randomised, 335 (58%) achieved clinical response by week 2. These patients were classified as week-2 responders and were included in the primary analysis (see Table 5). Among patients classified as non-responders at week 2, 32% (26/81) in the placebo maintenance group and 42% (68/163) in the infliximab group achieved clinical response by week 6. There was no difference between groups in the number of late responders thereafter.

The co-primary endpoints were the proportion of patients in clinical remission (CDAI  $< 150$ ) at week 30 and time to loss of response through week 54. Corticosteroid tapering was permitted after week 6.

**Table 5**  
**Effects on response and remission rate, data from ACCENT I (week-2 responders)**

	ACCENT I (week-2 responders) % of Patients		
	Placebo Maintenance (n=110)	Infliximab Maintenance 5 mg/kg (n=113) (p value)	Infliximab Maintenance 10 mg/kg (n=112) (p value)
Median time to loss of response through week 54	19 weeks	38 weeks (0.002)	> 54 weeks ( $< 0.001$ )
<b>Week 30</b>			
Clinical Response <sup>a</sup>	27.3	51.3 ( $< 0.001$ )	59.1 ( $< 0.001$ )
Clinical Remission	20.9	38.9 (0.003)	45.5 ( $< 0.001$ )
Steroid-Free Remission	10.7 (6/56)	31.0 (18/58) (0.008)	36.8 (21/57) (0.001)
<b>Week 54</b>			
Clinical Response <sup>a</sup>	15.5	38.1 ( $< 0.001$ )	47.7 ( $< 0.001$ )
Clinical Remission	13.6	28.3 (0.007)	38.4 ( $< 0.001$ )
Sustained Steroid-Free Remission <sup>b</sup>	5.7 (3/53)	17.9 (10/56) (0.075)	28.6 (16/56) (0.002)

<sup>a</sup> Reduction in CDAI  $\geq 25\%$  and  $\geq 70$  points.

<sup>b</sup> CDAI  $< 150$  at both week 30 and 54 and not receiving corticosteroids in the 3 months prior to week 54 among patients who were receiving corticosteroids at baseline.

Beginning at week 14, patients who had responded to treatment, but subsequently lost their clinical benefit, were allowed to cross over to a dose of infliximab 5 mg/kg higher than the dose to which they were originally randomised. Eighty nine percent (50/56) of patients who lost clinical response on infliximab 5 mg/kg maintenance therapy after week 14 responded to treatment with infliximab 10 mg/kg.

Improvements in quality of life measures, a reduction in disease-related hospitalisations and corticosteroid use were seen in the infliximab maintenance groups compared with the placebo maintenance group at weeks 30 and 54.

Infliximab with or without AZA was assessed in a randomised, double-blind, active comparator study (SONIC) of 508 adult patients with moderate to severe Crohn's disease (CDAI  $\geq 220 \leq 450$ ) who were naive to biologics and immunosuppressants and had a median disease duration of 2.3 years. At baseline 27.4% of patients were receiving systemic corticosteroids, 14.2% of patients were receiving budesonide, and 54.3% of patients were receiving 5-ASA compounds. Patients were randomised to receive AZA monotherapy, infliximab monotherapy, or infliximab plus AZA combination therapy. Infliximab was administered at a dose of 5 mg/kg at weeks 0, 2, 6, and then every 8 weeks. AZA was given at a dose of 2.5 mg/kg daily.

The primary endpoint of the study was corticosteroid-free clinical remission at week 26, defined as patients in clinical remission (CDAI of  $< 150$ ) who, for at least 3 weeks, had not taken oral systemic corticosteroids (prednisone or equivalent) or budesonide at a dose  $> 6$  mg/day. For results see Table 6. The proportions of patients with mucosal healing at week 26 were significantly greater in the infliximab plus AZA combination (43.9%,  $p < 0.001$ ) and infliximab monotherapy groups (30.1%,  $p=0.023$ ) compared to the AZA monotherapy group (16.5%).

**Table 6**  
**Percent of patients achieving corticosteroid-free clinical remission at week 26, SONIC**

	AZA Monotherapy	Infliximab Monotherapy	Infliximab + AZA Combination therapy
<b>Week 26</b>			
All randomised patients	30.0% (51/170)	44.4% (75/169) ( $p=0.006$ )*	56.8% (96/169) ( $p<0.001$ )*

\* P-values represent each infliximab treatment group vs. AZA monotherapy

Similar trends in the achievement of corticosteroid-free clinical remission were observed at week 50. Furthermore, improved quality of life as measured by IBDQ was observed with infliximab.

#### *Induction treatment in fistulising active Crohn's disease*

The efficacy was assessed in a randomised, double-blinded, placebo-controlled study in 94 patients with fistulising Crohn's disease who had fistulae that were of at least 3 months' duration. Thirty one of these patients were treated with infliximab 5 mg/kg. Approximately 93% of the patients had previously received antibiotic or immunosuppressive therapy.

Concurrent use of stable doses of conventional therapies was permitted, and 83% of patients continued to receive at least one of these therapies. Patients received three doses of either placebo or infliximab at weeks 0, 2 and 6. Patients were followed up to 26 weeks. The primary endpoint was the proportion of patients who experienced a clinical response, defined as  $\geq 50\%$  reduction from baseline in the number of fistulae draining upon gentle compression on at least two consecutive visits (4 weeks apart), without an increase in the use of medicinal products or surgery for Crohn's disease.

Sixty eight percent (21/31) of infliximab-treated patients receiving a 5 mg/kg dose regimen achieved a clinical response vs. 26% (8/31) placebo-treated patients ( $p=0.002$ ). The median time to onset of response in the infliximab-treated group was 2 weeks. The median duration of response was 12 weeks. Additionally, closure of all fistulae was achieved in 55% of infliximab-treated patients compared with 13% of placebo-treated patients ( $p=0.001$ ).

#### *Maintenance treatment in fistulising active Crohn's disease*

The efficacy of repeated infusions with infliximab in patients with fistulising Crohn's disease was studied in a 1-year clinical study (ACCENT II). A total of 306 patients received 3 doses of infliximab 5 mg/kg at week 0, 2 and 6. At baseline, 87% of the patients had perianal fistulae, 14% had abdominal fistulae, 9% had rectovaginal fistulae. The median CDAI score was 180. At week 14, 282 patients were assessed for clinical response and randomised to receive either placebo or 5 mg/kg infliximab every 8 weeks through week 46.

Week-14 responders (195/282) were analysed for the primary endpoint, which was time from randomisation to loss of response (see Table 7). Corticosteroid tapering was permitted after week 6.

**Table 7**  
**Effects on response rate, data from ACCENT II (week-14 responders)**

	ACCENT II (week-14 responders)		
	Placebo Maintenance (n=99)	Infliximab Maintenance (5 mg/kg) (n=96)	p-value
Median time to loss of response through week 54	14 weeks	> 40 weeks	< 0.001
<b>Week 54</b>			
Fistula Response (%) <sup>a</sup>	23.5	46.2	0.001
Complete fistula response (%) <sup>b</sup>	19.4	36.3	0.009

<sup>a</sup>  $\geq 50\%$  reduction from baseline in the number of draining fistulas over a period of  $\geq 4$  weeks

<sup>b</sup> Absence of any draining fistulas

Beginning at week 22, patients who initially responded to treatment and subsequently lost their response were eligible to cross over to active re-treatment every 8 weeks at a dose of infliximab 5 mg/kg higher than the dose to which they were originally randomised. Among patients in the infliximab 5 mg/kg group who crossed over because of loss of fistula response after week 22, 57% (12/21) responded to re-treatment with infliximab 10 mg/kg every 8 weeks.

There was no significant difference between placebo and infliximab for the proportion of patients with sustained closure of all fistulas through week 54, for symptoms such as proctalgia, abscesses and urinary tract infection or for number of newly developed fistulas during treatment.

Maintenance therapy with infliximab every 8 weeks significantly reduced disease-related hospitalisations and surgeries compared with placebo. Furthermore, a reduction in corticosteroid use and improvements in quality of life were observed.

#### Adult ulcerative colitis

The safety and efficacy of infliximab were assessed in two (ACT 1 and ACT 2) randomised, double-blind, placebo-controlled clinical studies in adult patients with moderately to severely active ulcerative colitis (Mayo score 6 to 12; Endoscopy subscore  $\geq 2$ ) with an inadequate response to conventional therapies [oral corticosteroids, aminosalicylates and/or immunomodulators (6-MP, AZA)]. Concomitant stable doses of oral aminosalicylates, corticosteroids, and/or immunomodulatory agents were permitted. In both studies, patients were randomised to receive either placebo, 5 mg/kg infliximab, or 10 mg/kg infliximab at weeks 0, 2, 6, 14 and 22, and in ACT 1 at weeks 30, 38 and 46. Corticosteroid taper was permitted after week 8.

**Table 8**  
**Effects on clinical response, clinical remission and mucosal healing at weeks 8 and 30.**  
**Combined data from ACT 1 & 2**

	Placebo	Infliximab		
		5 mg/kg	10 mg/kg	Combined
Subjects randomised	244	242	242	484
<b>Percentage of subjects in clinical response and in sustained clinical response</b>				
Clinical response at week 8 <sup>a</sup>	33.2%	66.9%	65.3%	66.1%
Clinical response at week 30 <sup>a</sup>	27.9%	49.6%	55.4%	52.5%
Sustained response (clinical response at both week 8 and week 30) <sup>a</sup>	19.3%	45.0%	49.6%	47.3%
<b>Percentage of subjects in clinical remission and sustained remission</b>				
Clinical remission at week 8 <sup>a</sup>	10.2%	36.4%	29.8%	33.1%
Clinical remission at week 30 <sup>a</sup>	13.1%	29.8%	36.4%	33.1%
Sustained remission (in remission at both week 8 and week 30) <sup>a</sup>	5.3%	19.0%	24.4%	21.7%
<b>Percentage of subjects with mucosal healing</b>				
Mucosal healing at week 8 <sup>a</sup>	32.4%	61.2%	60.3%	60.7%
Mucosal healing at week 30 <sup>a</sup>	27.5%	48.3%	52.9%	50.6%

<sup>a</sup>  $p < 0.001$ , for each infliximab treatment group vs. placebo

The efficacy of infliximab through week 54 was assessed in the ACT 1 study.

At 54 weeks, 44.9% of patients in the combined infliximab treatment group were in clinical response compared to 19.8% in the placebo treatment group ( $p < 0.001$ ). Clinical remission and mucosal healing occurred in a greater proportion of patients in the combined infliximab treatment group compared to the placebo treatment group at week 54 (34.6% vs. 16.5%,  $p < 0.001$  and 46.1% vs. 18.2%,  $p < 0.001$ , respectively). The proportions of patients in sustained response and sustained remission at week 54 were greater in the combined infliximab treatment group than in the placebo treatment group (37.9% vs. 14.0%,  $p < 0.001$ ; and 20.2% vs. 6.6%,  $p < 0.001$ , respectively).

A greater proportion of patients in the combined infliximab treatment group were able to discontinue corticosteroids while remaining in clinical remission compared to the placebo treatment group at both week 30 (22.3% vs. 7.2%,  $p < 0.001$ , pooled ACT 1 & ACT 2 data) and week 54 (21.0% vs. 8.9%,  $p=0.022$ , ACT 1 data).

The pooled data analysis from the ACT 1 and ACT 2 studies and their extensions, analysed from baseline through 54 weeks, demonstrated a reduction of ulcerative colitis-related hospitalisations and surgical

procedures with infliximab treatment. The number of ulcerative colitis-related hospitalisations was significantly lower in the 5 and 10 mg/kg infliximab treatment groups than in the placebo group (mean number of hospitalisations per 100 subject-years: 21 and 19 vs. 40 in the placebo group;  $p=0.019$  and  $p=0.007$ , respectively). The number of ulcerative colitis-related surgical procedures was also lower in the 5 and 10 mg/kg infliximab treatment groups than in the placebo group (mean number of surgical procedures per 100 subject-years: 22 and 19 vs. 34;  $p=0.145$  and  $p=0.022$ , respectively).

The proportion of subjects who underwent colectomy at any time within 54 weeks following the first infusion of study agent were collected and pooled from the ACT 1 and ACT 2 studies and their extensions. Fewer subjects underwent colectomy in the 5 mg/kg infliximab group (28/242 or 11.6% [N.S.]) and the 10 mg/kg infliximab group (18/242 or 7.4% [ $p=0.011$ ]) than in the placebo group (36/244; 14.8%).

The reduction in incidence of colectomy was also examined in another randomised, double-blind study (C0168Y06) in hospitalised patients ( $n=45$ ) with moderately to severely active ulcerative colitis who failed to respond to intravenous (IV) corticosteroids and who were therefore at higher risk for colectomy. Significantly fewer colectomies occurred within 3 months of study infusion in patients who received a single dose of 5 mg/kg infliximab compared to patients who received placebo (29.2% vs. 66.7% respectively,  $p=0.017$ ).

In ACT 1 and ACT 2, infliximab improved quality of life, confirmed by statistically significant improvement in both a disease specific measure, IBDQ, and by improvement in the generic 36-item short form survey SF-36.

#### Adult ankylosing spondylitis

Efficacy and safety of infliximab were assessed in two multicentre, double-blind, placebo-controlled studies in patients with active ankylosing spondylitis (Bath Ankylosing Spondylitis Disease Activity Index [BASDAI] score  $\geq 4$  and spinal pain  $\geq 4$  on a scale of 1-10).

In the first study (P01522), which had a 3 month double-blind phase, 70 patients received either 5 mg/kg infliximab or placebo at weeks 0, 2, 6 (35 patients in each group). At week 12, placebo patients were switched to infliximab 5 mg/kg every 6 weeks up to week 54. After the first year of the study, 53 patients continued into an open-label extension to week 102.

In the second clinical study (ASSERT), 279 patients were randomised to receive either placebo (Group 1,  $n=78$ ) or 5 mg/kg infliximab (Group 2,  $n=201$ ) at 0, 2 and 6 weeks and every 6 weeks to week 24. Thereafter, all subjects continued on infliximab every 6 weeks to week 96. Group 1 received 5 mg/kg infliximab. In Group 2, starting with the week 36 infusion, patients who had a BASDAI  $\geq 3$  at 2 consecutive visits, received 7.5 mg/kg infliximab every 6 weeks thereafter through week 96.

In ASSERT, improvement in signs and symptoms was observed as early as week 2. At week 24, the number of ASAS 20 responders was 15/78 (19%) in the placebo group, and 123/201 (61%) in the 5 mg/kg infliximab group ( $p < 0.001$ ). There were 95 subjects from group 2 who continued on 5 mg/kg every 6 weeks. At 102 weeks there were 80 subjects still on infliximab treatment and among those, 71 (89%) were ASAS 20 responders.

In P01522, improvement in signs and symptoms was also observed as early as week 2. At week 12, the number of BASDAI 50 responders were 3/35 (9%) in the placebo group, and 20/35 (57%) in the 5 mg/kg group ( $p < 0.01$ ). There were 53 subjects who continued on 5 mg/kg every 6 weeks. At 102 weeks there were 49 subjects still on infliximab treatment and among those, 30 (61%) were BASDAI 50 responders.

In both studies, physical function and quality of life as measured by the BASFI and the physical component score of the SF-36 were also improved significantly.

#### Adult psoriatic arthritis

Efficacy and safety were assessed in two multicentre, double-blind, placebo-controlled studies in patients with active psoriatic arthritis.

In the first clinical study (IMPACT), efficacy and safety of infliximab were studied in 104 patients with active polyarticular psoriatic arthritis. During the 16-week double-blind phase, patients received either 5 mg/kg infliximab or placebo at weeks 0, 2, 6, and 14 (52 patients in each group). Starting at week 16, placebo patients were switched to infliximab and all patients subsequently received 5 mg/kg infliximab every 8 weeks up to week 46. After the first year of the study, 78 patients continued into an open-label extension to week 98.

In the second clinical study (IMPACT 2), efficacy and safety of infliximab were studied in 200 patients with active psoriatic arthritis ( $\geq 5$  swollen joints and  $\geq 5$  tender joints). Forty six percent of patients continued on stable doses of methotrexate ( $\leq 25$  mg/week). During the 24-week double-blind phase, patients received either 5 mg/kg infliximab or placebo at weeks 0, 2, 6, 14, and 22 (100 patients in each group). At week 16, 47 placebo patients with  $< 10\%$  improvement from baseline in both swollen and tender joint counts were switched to infliximab induction (early escape). At week 24, all placebo-treated patients crossed over to infliximab induction. Dosing continued for all patients through week 46.

Key efficacy results for IMPACT and IMPACT 2 are shown in Table 9 below:

**Table 9**  
**Effects on ACR and PASI in IMPACT and IMPACT 2**

	IMPACT			IMPACT 2*		
	Placebo (week 16)	Infliximab (week 16)	Infliximab (week 98)	Placebo (week 24)	Infliximab (week 24)	Infliximab (week 54)
Patients randomised	52	52	N/A <sup>a</sup>	100	100	100
ACR response (% of patients)						
N	52	52	78	100	100	100
ACR 20 response*	5(10%)	34 (65%)	48 (62%)	16 (16%)	54 (54%)	53 (53%)
ACR 50 response*	0(0%)	24 (46%)	35 (45%)	4 (4%)	41(41%)	33 (33%)
ACR 70 response*	0(0%)	15 (29%)	27 (35%)	2 (2%)	27 (27%)	20 (20%)
PASI response (% of patients) <sup>b</sup>						
N	N/A			87	83	82
PASI 75 response**	N/A			1 (1%)	50 (60%)	40 (48.8%)

\* ITT-analysis where subjects with missing data were included as non-responders

<sup>a</sup> Week 98 data for IMPACT includes combined placebo crossover and infliximab patients who entered the open-label extension

<sup>b</sup> Based on patients with PASI  $\geq 2.5$  at baseline for IMPACT, and patients with  $\geq 3\%$  BSA psoriasis skin involvement at baseline in IMPACT 2

\*\* PASI 75 response for IMPACT not included due to low N;  $p < 0.001$  for infliximab vs. placebo at week 24 for IMPACT 2

In IMPACT and IMPACT 2, clinical responses were observed as early as week 2 and were maintained through week 98 and week 54 respectively. Efficacy has been demonstrated with or without concomitant use of methotrexate. Decreases in parameters of peripheral activity characteristic of psoriatic arthritis (such as number of swollen joints, number of painful/tender joints, dactylitis and presence of enthesopathy) were seen in the infliximab-treated patients.

Radiographic changes were assessed in IMPACT 2. Radiographs of hands and feet were collected at baseline, weeks 24 and 54. Infliximab treatment reduced the rate of progression of peripheral joint damage compared with placebo treatment at the week 24 primary endpoint as measured by change from baseline in total modified vdH-S score (mean  $\pm$  SD score was  $0.82 \pm 2.62$  in the placebo group compared with  $-0.70 \pm 2.53$  in the infliximab group;  $p < 0.001$ ). In the infliximab group, the mean change in total modified vdH-S score remained below 0 at the week 54 timepoint.

Infliximab-treated patients demonstrated significant improvement in physical function as assessed by HAQ. Significant improvements in health-related quality of life were also demonstrated as measured by the physical and mental component summary scores of the SF-36 in IMPACT 2.

### Adult psoriasis

The efficacy of infliximab was assessed in two multicentre, randomised, double-blind studies: SPIRIT and EXPRESS. Patients in both studies had plaque psoriasis (Body Surface Area [BSA]  $\geq$  10% and Psoriasis Area and Severity Index [PASI] score  $\geq$  12). The primary endpoint in both studies was the percent of patients who achieved  $\geq$  75% improvement in PASI from baseline at week 10.

SPIRIT evaluated the efficacy of infliximab induction therapy in 249 patients with plaque psoriasis that had previously received PUVA or systemic therapy. Patients received either 3 or 5 mg/kg infliximab or placebo infusions at weeks 0, 2 and 6. Patients with a PGA score  $\geq$  3 were eligible to receive an additional infusion of the same treatment at week 26.

In SPIRIT, the proportion of patients achieving PASI 75 at week 10 was 71.7% in the 3 mg/kg infliximab group, 87.9% in the 5 mg/kg infliximab group, and 5.9% in the placebo group ( $p < 0.001$ ). By week 26, twenty weeks after the last induction dose, 30% of patients in the 5 mg/kg group and 13.8% of patients in the 3 mg/kg group were PASI 75 responders. Between weeks 6 and 26, symptoms of psoriasis gradually returned with a median time to disease relapse of  $> 20$  weeks. No rebound was observed.

EXPRESS evaluated the efficacy of infliximab induction and maintenance therapy in 378 patients with plaque psoriasis. Patients received 5 mg/kg infliximab- or placebo-infusions at weeks 0, 2 and 6 followed by maintenance therapy every 8 weeks through week 22 in the placebo group and through week 46 in the infliximab group. At week 24, the placebo group crossed over to infliximab induction therapy (5 mg/kg) followed by infliximab maintenance therapy (5 mg/kg). Nail psoriasis was assessed using the Nail Psoriasis Severity Index (NAPSI). Prior therapy with PUVA, methotrexate, cyclosporin, or acitretin had been received by 71.4% of patients, although they were not necessarily therapy resistant. Key results are presented in Table 10. In infliximab treated subjects, significant PASI 50 responses were apparent at the first visit (week 2) and PASI 75 responses by the second visit (week 6). Efficacy was similar in the subgroup of patients that were exposed to previous systemic therapies compared to the overall study population.

**Table 10**  
**Summary of PASI response, PGA response and percent of patients with all nails cleared at weeks 10, 24 and 50. EXPRESS**

	Placebo $\rightarrow$ Infliximab 5 mg/kg (at week 24)	Infliximab 5 mg/kg
<b>Week 10</b>		
N	77	301
$\geq$ 90% improvement	1 (1.3%)	172 (57.1%) <sup>a</sup>
$\geq$ 75% improvement	2 (2.6%)	242 (80.4%) <sup>a</sup>
$\geq$ 50% improvement	6 (7.8%)	274 (91.0%)
PGA of cleared (0) or minimal (1)	3 (3.9%)	242 (82.9%) <sup>ab</sup>
PGA of cleared (0), minimal (1), or mild (2)	14 (18.2%)	275 (94.2%) <sup>ab</sup>
<b>Week 24</b>		
N	77	276
$\geq$ 90% improvement	1 (1.3%)	161 (58.3%) <sup>a</sup>
$\geq$ 75% improvement	3 (3.9%)	227 (82.2%) <sup>a</sup>
$\geq$ 50% improvement	5 (6.5%)	248 (89.9%)
PGA of cleared (0) or minimal (1)	2 (2.6%)	203 (73.6%) <sup>a</sup>
PGA of cleared (0), minimal (1), or mild (2)	15 (19.5%)	246 (89.1%) <sup>a</sup>
<b>Week 50</b>		
N	68	281
$\geq$ 90% improvement	34 (50.0%)	127 (45.2%)
$\geq$ 75% improvement	52 (76.5%)	170 (60.5%)
$\geq$ 50% improvement	61 (89.7%)	193 (68.7%)
PGA of cleared (0) or minimal (1)	46 (67.6%)	149 (53.0%)
PGA of cleared (0), minimal (1), or mild (2)	59 (86.8%)	189 (67.3%)
<b>All nails cleared<sup>c</sup></b>		
Week 10	1/65 (1.5%)	16/235 (6.8%)

	Placebo → Infliximab 5 mg/kg (at week 24)	Infliximab 5 mg/kg
Week 24	3/65 (4.6%)	58/223 (26.0%) <sup>a</sup>
Week 50	27/64 (42.2%)	92/226 (40.7%)

<sup>a</sup> p < 0.001, for each infliximab treatment group vs. control

<sup>b</sup> n = 292

<sup>c</sup> Analysis was based on subjects with nail psoriasis at baseline (81.8% of subjects). Mean baseline NAPSI scores were 4.6 and 4.3 in infliximab and placebo group.

Significant improvements from baseline were demonstrated in DLQI (p < 0.001) and the physical and mental component scores of the SF 36 (p < 0.001 for each component comparison).

### Paediatric population

#### Paediatric Crohn's disease (6 to 17 years)

In the REACH study, 112 patients (6 to 17 years, median age 13.0 years) with moderate to severe, active Crohn's disease (median paediatric CDAI of 40) and an inadequate response to conventional therapies were to receive 5 mg/kg infliximab at weeks 0, 2, and 6. All patients were required to be on a stable dose of 6-MP, AZA or MTX (35% were also receiving corticosteroids at baseline). Patients assessed by the investigator to be in clinical response at week 10 were randomised and received 5 mg/kg infliximab at either q8 weeks or q12 weeks as a maintenance treatment regimen. If response was lost during maintenance treatment, crossing over to a higher dose (10 mg/kg) and/or shorter dosing interval (q8 weeks) was allowed. Thirty two (32) evaluable paediatric patients crossed over (9 subjects in the q8 weeks and 23 subjects in the q12 weeks maintenance groups). Twenty four of these patients (75.0%) regained clinical response after crossing over. The proportion of subjects in clinical response at week 10 was 88.4% (99/112). The proportion of subjects achieving clinical remission at week 10 was 58.9% (66/112).

At week 30, the proportion of subjects in clinical remission was higher in the q8 week (59.6%, 31/52) than the q12 week maintenance treatment group (35.3%, 18/51; p=0.013). At week 54, the figures were 55.8% (29/52) and 23.5% (12/51) in the q8 weeks and q12 weeks maintenance groups, respectively (p < 0.001). Data about fistulas were derived from PCDAI scores. Of the 22 subjects that had fistulas at baseline, 63.6% (14/22), 59.1% (13/22) and 68.2% (15/22) were in complete fistula response at week 10, 30 and 54, respectively, in the combined q8 weeks and q12 weeks maintenance groups.

In addition, statistically and clinically significant improvements in quality of life and height, as well as a significant reduction in corticosteroid use, were observed versus baseline.

#### Paediatric ulcerative colitis (6 to 17 years)

The safety and efficacy of infliximab were assessed in a multicentre, randomised, open-label, parallel-group clinical study (C0168T72) in 60 paediatric patients aged 6 through 17 years (median age 14.5 years) with moderately to severely active ulcerative colitis (Mayo score of 6 to 12; endoscopic subscore ≥ 2) with an inadequate response to conventional therapies. At baseline 53% of patients were receiving immunomodulator therapy (6-MP, AZA and/or MTX) and 62% of patients were receiving corticosteroids. Discontinuation of immunomodulators and corticosteroid taper were permitted after week 0.

All patients received an induction regimen of 5 mg/kg infliximab at weeks 0, 2, and 6. Patients who did not respond to infliximab at week 8 (n=15) received no further medicinal product and returned for safety follow-up. At week 8, 45 patients were randomised and received 5 mg/kg infliximab at either q8 weeks or q12 weeks as a maintenance treatment regimen.

The proportion of patients in clinical response at week 8 was 73.3% (44/60). Clinical response at week 8 was similar between those with or without concomitant immunomodulator use at baseline. Clinical remission at week 8 was 33.3% (17/51) as measured by the Paediatric Ulcerative Colitis Activity Index (PUCAI) score.

At week 54, the proportion of patients in clinical remission as measured by the PUCAI score was 38% (8/21) in the q8 week maintenance group and 18% (4/22) in the q12 week maintenance treatment group. For patients receiving corticosteroids at baseline, the proportion of patients in remission and not receiving corticosteroids at week 54 was 38.5% (5/13) for the q8 week and 0% (0/13) for the q12 week maintenance treatment group.

In this study, there were more patients in the 12 to 17 year age group than in the 6 to 11 year age group (45/60 vs.15/60). While the numbers of patients in each subgroup are too small to draw definitive conclusions about the effect of age, there was a higher number of patients in the younger age group who stepped up in dose or discontinued treatment due to inadequate efficacy.

#### Other paediatric indications

The European Medicines Agency has waived the obligation to submit the results of studies with the reference medicinal product containing infliximab in all subsets of the paediatric population in rheumatoid arthritis, juvenile idiopathic arthritis, psoriatic arthritis, ankylosing spondylitis, psoriasis and Crohn's disease (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

Single intravenous infusions of 1, 3, 5, 10 or 20 mg/kg of infliximab yielded dose proportional increases in the maximum serum concentration ( $C_{max}$ ) and area under the concentration-time curve (AUC). The volume of distribution at steady state (median  $V_d$  of 3.0 to 4.1 litres) was not dependent on the administered dose and indicated that infliximab is predominantly distributed within the vascular compartment. No time-dependency of the Pharmacokinetics was observed. The elimination pathways for infliximab have not been characterised. Unchanged infliximab was not detected in urine. No major age- or weight-related differences in clearance or volume of distribution were observed in rheumatoid arthritis patients. The pharmacokinetics of infliximab in elderly patients has not been studied. Studies have not been performed in patients with liver or renal disease.

At single doses of 3, 5, or 10 mg/kg, the median  $C_{max}$  values were 77, 118 and 277 micrograms/mL, respectively. The median terminal half-life at these doses ranged from 8 to 9.5 days. In most patients, infliximab could be detected in the serum for at least 8 weeks after the recommended single dose of 5 mg/kg for Crohn's disease and the rheumatoid arthritis maintenance dose of 3 mg/kg every 8 weeks.

Repeated administration of infliximab (5 mg/kg at 0, 2 and 6 weeks in fistulising Crohn's disease, 3 or 10 mg/kg every 4 or 8 weeks in rheumatoid arthritis) resulted in a slight accumulation of infliximab in serum after the second dose. No further clinically relevant accumulation was observed. In most fistulising Crohn's disease patients, infliximab was detected in serum for 12 weeks (range 4-28 weeks) after administration of the regimen.

#### *Paediatric population*

Population pharmacokinetic analysis based on data obtained from patients with ulcerative colitis (N=60), Crohn's disease (N=112), juvenile rheumatoid arthritis (N=117) and Kawasaki disease (N=16) with an overall age range from 2 months to 17 years indicated that exposure to infliximab was dependent on body weight in a non-linear way. Following administration of 5 mg/kg infliximab every 8 weeks, the predicted median steady-state infliximab exposure (area under concentration-time curve at steady state,  $AUC_{ss}$ ) in paediatric patients aged 6 years to 17 years was approximately 20% lower than the predicted median steady-state drug exposure in adults. The median  $AUC_{ss}$  in paediatric patients aged 2 years to less than 6 years was predicted to be approximately 40% lower than that in adults, although the number of patients supporting this estimate is limited.

## **5.3 Preclinical safety data**

Infliximab does not cross react with  $TNF_{\alpha}$  from species other than human and chimpanzees. Therefore, conventional preclinical safety data with infliximab are limited. In a developmental toxicity study conducted in mice using an analogous antibody that selectively inhibits the functional activity of mouse  $TNF_{\alpha}$ , there was no indication of maternal toxicity, embryotoxicity or teratogenicity. In a fertility and general reproductive function study, the number of pregnant mice was reduced following administration of the same analogous antibody. It is not known whether this finding was due to effects on the males and/or the females. In a 6-month repeated dose toxicity study in mice, using the same analogous antibody against mouse  $TNF_{\alpha}$ , crystalline deposits were observed on the lens capsule of some of the treated male mice. No specific

ophthalmologic examinations have been performed in patients to investigate the relevance of this finding for humans.

Long-term studies have not been performed to evaluate the carcinogenic potential of infliximab. Studies in mice deficient in TNF $\alpha$  demonstrated no increase in tumours when challenged with known tumour initiators and/or promoters.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Sucrose

Polysorbate 80

Monobasic sodium phosphate monohydrate

Dibasic sodium phosphate heptahydrate

### **6.2 Incompatibilities**

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

36 months.

Chemical and physical in use stability of the reconstituted solution has been demonstrated for 24 hours at 25°C. From a microbiological point of view, the product should be used as soon as possible but within 3 hours of reconstitution and dilution. If not used immediately, in use storage times and conditions prior to use are the responsibility of the user and should not be longer than 24 hours at 2°C to 8°C.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C).

For storage conditions after reconstitution of the medicinal product, see section 6.3.

### **6.5 Nature and contents of container**

Type 1 glass vial with a rubber stopper and aluminium crimp protected by a plastic cap, containing 100 mg of infliximab.

Flixabi is available in packs containing 1 vial and multipacks containing 2, 3, 4, and 5 (2, 3, 4, and 5 packs of 1) vials. Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal and other handling**

1. Calculate the dose and the number of Flixabi vials needed. Each Flixabi vial contains 100 mg infliximab. Calculate the total volume of reconstituted Flixabi solution required.
2. Under aseptic conditions, reconstitute each Flixabi vial with 10 ml of water for injections, using a syringe equipped with a 21-gauge (0.8 mm) or smaller needle. Remove flip-top from the vial and wipe the top with a 70% alcohol swab. Insert the syringe needle into the vial through the centre of the rubber stopper and direct the stream of water for injections to the glass wall of the vial. Gently swirl the solution by rotating the vial to dissolve the lyophilised powder. Avoid prolonged or vigorous agitation. **DO NOT SHAKE**. Foaming of the solution on reconstitution is not unusual. Allow the reconstituted

solution to stand for 5 minutes. Check that the solution is colourless to light yellow and opalescent. The solution may develop a few fine translucent particles, as infliximab is a protein. Do not use if opaque particles, discolouration, or other foreign particles are present.

3. Dilute the total volume of the reconstituted Flixabi solution dose to 250 ml with sodium chloride 9 mg/ml (0.9%) solution for infusion. Do not dilute the reconstituted Flixabi solution with any other diluent. The dilution can be accomplished by withdrawing a volume of the sodium chloride 9 mg/ml (0.9%) solution for infusion from the 250-ml glass bottle or infusion bag equal to the volume of reconstituted Flixabi. Slowly add the total volume of reconstituted Flixabi solution to the 250-ml infusion bottle or bag. Gently mix.
4. Administer the infusion solution over a period of not less than the infusion time recommended (see section 4.2). Use only an infusion set with an in-line, sterile, non-pyrogenic, low protein-binding filter (pore size 1.2 micrometre or less). Since no preservative is present, it is recommended that the administration of the solution for infusion is to be started as soon as possible and within 3 hours of reconstitution and dilution. When reconstitution and dilution are performed under aseptic conditions, Flixabi infusion solution can be used within 24 hours if stored at 2°C to 8°C. Do not store any unused portion of the infusion solution for reuse.
5. No physical biochemical compatibility studies have been conducted to evaluate the co-administration of Flixabi with other agents. Do not infuse Flixabi concomitantly in the same intravenous line with other agents.
6. Visually inspect Flixabi for particulate matter or discolouration prior to administration. Do not use if visibly opaque particles, discolouration or foreign particles are observed.
7. Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Samsung Bioepis UK Limited  
5th Floor, Profile West, 950 Great West Road  
Brentford  
Middlesex TW8 9ES  
United Kingdom

## **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/16/1106/001  
EU/1/16/1106/002  
EU/1/16/1106/003  
EU/1/16/1106/004  
EU/1/16/1106/005

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 26/05/2016

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the website of the European Medicines Agency  
<http://www.ema.europa.eu>

## **ANNEX II**

- A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

### Name and address of the manufacturer of the biological active substance

Biogen (Denmark) Manufacturing ApS  
Biogen Alle' 1  
3400 Hillerod  
DENMARK

### Name and address of the manufacturer responsible for batch release

Biogen (Denmark) Manufacturing ApS  
Biogen Alle' 1  
3400 Hillerod  
DENMARK

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports**

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

- **Additional risk minimisation measures**

Prior to launch in each Member State, the MAH shall agree the final educational material with the competent authority in that Member State, consisting of information provided to all healthcare professionals expected to prescribe the product.

The healthcare professional's educational material should contain the following key elements:

- The risk of opportunistic infections and tuberculosis (TB) in patients treated with Flixabi.
- The need to assess the risk of TB in patients prior to treating with Flixabi.
- The risk of acute hypersensitivity reactions (including anaphylactic shock) and delayed hypersensitivity reactions.
- The risk of lymphoma, melanoma, Merkel cell carcinoma, and other malignancies.
- The risk of disseminated BCG infection after BCG vaccination of infants up to 6 months of age who were exposed in utero to infliximab.
- The patient alert card, which is to be given to patients using Flixabi.

Prescribers of Flixabi for paediatric Crohn's disease and paediatric ulcerative colitis shall additionally be made aware:

- That children may be at increased risk of developing infections and that their immunisations need to be up-to-date.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Flixabi 100 mg powder for concentrate for solution for infusion  
Infliximab

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each vial contains 100 mg of infliximab.  
After reconstitution each mL contains 10 mg infliximab.

**3. LIST OF EXCIPIENTS**

Excipients: sucrose, polysorbate 80, monobasic sodium phosphate monohydrate and dibasic sodium phosphate heptahydrate.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Powder for concentrate for solution for infusion  
1 vial

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For intravenous use after reconstitution and dilution.  
Read the package leaflet before reconstitution and use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP:

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Samsung Bioepis UK Limited  
5th Floor, Profile West, 950 Great West Road  
Brentford  
Middlesex TW8 9ES  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/16/1106/001

**13. BATCH NUMBER**

Lot:

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

<2D barcode carrying the unique identifier included.>

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON FOR MULTIPACKS OF 2, 3, 4, AND 5 VIALS (WITH BLUE BOX)**

**1. NAME OF THE MEDICINAL PRODUCT**

Flixabi 100 mg powder for concentrate for solution for infusion  
Infliximab

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each vial contains 100 mg of infliximab.  
After reconstitution each mL contains 10 mg infliximab.

**3. LIST OF EXCIPIENTS**

Excipients: sucrose, polysorbate 80, monobasic sodium phosphate monohydrate and dibasic sodium phosphate heptahydrate.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Powder for concentrate for solution for infusion

Multipack: 2 (2 packs of 1) vials

Multipack: 3 (3 packs of 1) vials

Multipack: 4 (4 packs of 1) vials

Multipack: 5 (5 packs of 1) vials

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For intravenous use after reconstitution and dilution.  
Read the package leaflet before reconstitution and use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP:

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Samsung Bioepis UK Limited  
5th Floor, Profile West, 950 Great West Road  
Brentford  
Middlesex TW8 9ES  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/16/1106/002 multipack of 2 vials (2 packs of 1 vial)  
EU/1/16/1106/003 multipack of 3 vials (3 packs of 1 vial)  
EU/1/16/1106/004 multipack of 4 vials (4 packs of 1 vial)  
EU/1/16/1106/005 multipack of 5 vials (5 packs of 1 vial)

**13. BATCH NUMBER**

Lot:

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

<2D barcode carrying the unique identifier included.>

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:

SN:

NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON FOR 1 VIAL AS INTERMEDIATE PACK, COMPONENT OF A MULTIPACK  
(WITHOUT BLUE BOX)**

**1. NAME OF THE MEDICINAL PRODUCT**

Flixabi 100 mg powder for concentrate for solution for infusion  
Infliximab

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each vial contains 100 mg of infliximab.  
After reconstitution each mL contains 10 mg infliximab.

**3. LIST OF EXCIPIENTS**

Excipients: sucrose, polysorbate 80, monobasic sodium phosphate monohydrate and dibasic sodium phosphate heptahydrate.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Powder for concentrate for solution for infusion  
1 vial

Component of a multipack, cannot be sold separately.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For intravenous use after reconstitution and dilution.  
Read the package leaflet before reconstitution and use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF  
THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP:

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Samsung Bioepis UK Limited  
5th Floor, Profile West, 950 Great West Road  
Brentford  
Middlesex TW8 9ES  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/16/1106/002 multipack of 2 vials (2 packs of 1 vial)

EU/1/16/1106/003 multipack of 3 vials (3 packs of 1 vial)

EU/1/16/1106/004 multipack of 4 vials (4 packs of 1 vial)

EU/1/16/1106/005 multipack of 5 vials (5 packs of 1 vial)

**13. BATCH NUMBER**

Lot:

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

<2D barcode carrying the unique identifier included.>

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**VIAL LABEL**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Flixabi 100 mg powder for concentrate for solution for infusion  
Infliximab  
Intravenous use <IV>

**2. METHOD OF ADMINISTRATION**

**3. EXPIRY DATE**

EXP:

**4. BATCH NUMBER**

Lot:

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

100 mg

**6. OTHER**

## Flixabi 100 mg

Infliximab

### Patient Alert Card

Name patient:

Name doctor:

Telephone number doctor:

This Alert Card contains important safety information that you need to be aware of before and during treatment with Flixabi.

Show this card to any doctor involved in your treatment.

Please read the Flixabi 'Package Leaflet' carefully before you start using this medicine.

Date of Flixabi therapy initiation:

Current administrations:

It is important that you and your doctor record the brand name and batch number of your medicines.

Ask your doctor to record the type and date of last screening(s) for TB below:

Test:

Date:

Result:

Please make sure you also have a list of all other medicines that you are using with you at any visit to a healthcare professional.

List of allergies:

List of other medicines:

### Infections

#### Before treatment with Flixabi

- Tell your doctor if you have an infection even if it is a very minor one.
- It is very important that you tell your doctor if you have ever had tuberculosis (TB), or if you have been in close contact with someone who has had TB. Your doctor will test you to see if you have TB. Ask your doctor to record the type and date of your last screening(s) for TB on the card.
- Tell your doctor if you have hepatitis B or if you know or suspect you are a carrier of the hepatitis B virus.

#### During treatment with Flixabi

- Tell your doctor straight away if you have signs of an infection. Signs include a fever, feeling tired, (persistent) cough, shortness of breath, weight loss, night sweats, diarrhea, wounds, dental problems, burning when urinating or 'flu-like' signs.

### Heart Failure

#### Before treatment with Flixabi

- Tell your doctor if you have any heart problems such as mild heart failure.

#### During treatment with Flixabi

- Tell your doctor straight away if you notice signs of a heart problem. Signs include shortness of breath, swelling of the feet or changes in your heartbeat.

### Pregnancy and Vaccinations

- In case you have received Flixabi while you were pregnant, it is important that you inform your baby's doctor about it before your baby receives any vaccine. Your baby should not receive a 'live vaccine', such as BCG (used to prevent tuberculosis) within 6 months after birth.

Keep this card with you for 4 months after your dose of Flixabi, or in case of pregnancy for at least 6 months after the birth of your baby. Side effects may occur a long time after your last dose.

## **B. PACKAGE LEAFLET**

## Package leaflet: Information for the user

### Flixabi 100 mg powder for concentrate for solution for infusion Infliximab

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- Your doctor will also give you a Patient Alert Card, which contains important safety information you need to be aware of before and during your treatment with Flixabi.
- If you have any further questions, ask your doctor.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Flixabi is and what it is used for
2. What you need to know before you use Flixabi
3. How Flixabi will be given
4. Possible side effects
5. How to store Flixabi
6. Contents of the pack and other information

#### 1. What Flixabi is and what it is used for

Flixabi contains the active substance called infliximab. Infliximab is a type of protein of human and mouse origin.

Flixabi belongs to a group of medicines called ‘TNF blockers’. It is used in adults for the following inflammatory diseases:

- Rheumatoid arthritis
- Psoriatic arthritis
- Ankylosing spondylitis (Bechterew’s disease)
- Psoriasis.

Flixabi is also used in adults and children 6 years of age or older for:

- Crohn’s disease
- Ulcerative colitis.

Flixabi works by blocking the action of a protein called ‘tumour necrosis factor alpha’ (TNF<sub>α</sub>). This protein is involved in inflammatory processes of the body and blocking it can reduce the inflammation in your body.

#### Rheumatoid arthritis

Rheumatoid arthritis is an inflammatory disease of the joints. If you have active rheumatoid arthritis you will first be given other medicines. If you do not respond well enough to these medicines, you will be given Flixabi which you will take in combination with another medicine called methotrexate to:

- Reduce the signs and symptoms of your disease,
- Slow down the damage in your joints,

- Improve your physical function.

### **Psoriatic arthritis**

Psoriatic arthritis is an inflammatory disease of the joints, usually accompanied by psoriasis. If you have active psoriatic arthritis you will first be given other medicines. If you do not respond well enough to these medicines, you will be given Flixabi to:

- Reduce the signs and symptoms of your disease,
- Slow down the damage in your joints,
- Improve your physical function.

### **Ankylosing spondylitis (Bechterew's disease)**

Ankylosing spondylitis is an inflammatory disease of the spine. If you have ankylosing spondylitis you will first be given other medicines. If you do not respond well enough to these medicines, you will be given Flixabi to:

- Reduce the signs and symptoms of your disease,
- Improve your physical function.

### **Psoriasis**

Psoriasis is an inflammatory disease of the skin. If you have moderate to severe plaque psoriasis, you will first be given other medicines or treatments, such as phototherapy. If you do not respond well enough to these medicines or treatments, you will be given Flixabi to reduce the signs and symptoms of your disease.

### **Ulcerative colitis**

Ulcerative colitis is an inflammatory disease of the bowel. If you have ulcerative colitis you will first be given other medicines. If you do not respond well enough to these medicines, you will be given Flixabi to treat your disease.

### **Crohn's disease**

Crohn's disease is an inflammatory disease of the bowel. If you have Crohn's disease you will first be given other medicines. If you do not respond well enough to these, you will be given Flixabi to:

- Treat active Crohn's disease,
- Reduce the number of abnormal openings (fistulae) between your bowel and your skin that have not been controlled by other medicines or surgery.

## **2. What you need to know before you use Flixabi**

### **You should not be given Flixabi if**

- You are allergic to infliximab (the active ingredient in Flixabi) or any of the other ingredients of this medicine (listed in section 6).
- You are allergic (hypersensitive) to proteins that come from mice.
- You have tuberculosis (TB) or another serious infection such as pneumonia or sepsis.
- You have heart failure that is moderate or severe.

Do not have Flixabi if any of the above applies to you. If you are not sure, talk to your doctor before you are given Flixabi.

### **Warnings and precautions**

Talk to your doctor before you are given Flixabi if you have:

#### Had treatment with any medicine containing infliximab before

- Tell your doctor if you have had treatment with medicines containing infliximab in the past and are now starting Flixabi treatment again.
- If you have had a break in your treatment with infliximab of more than 16 weeks, there is a higher risk for allergic reactions when you start the treatment again.

### Infections

- Tell your doctor if you have an infection even if it is a very minor one before you are given Flixabi.
- Tell your doctor if you have lived in or travelled to an area where infections called histoplasmosis, coccidioidomycosis, or blastomycosis are common before you are given Flixabi. These infections are caused by specific types of fungi that can affect the lungs or other parts of your body.
- You may get infections more easily when you are being treated with Flixabi. If you are 65 or older, you have a greater risk.
- These infections may be serious and include tuberculosis, infections caused by viruses, fungi or bacteria, or other opportunistic infections and sepsis that may be life-threatening.

Tell your doctor straight away if you get signs of infection during treatment with Flixabi. Signs include fever, cough, flu-like signs, feeling unwell, red or hot skin, wounds or dental problems. Your doctor may recommend temporary discontinuation of Flixabi.

### Tuberculosis (TB)

- It is very important that you tell your doctor if you have ever had TB or if you have been in close contact with someone who has had or has TB.
- Your doctor will test you to see if you have TB. Cases of TB have been reported in patients treated with Flixabi, even in patients who have been treated with medicines for TB. Your doctor will record these tests on your Patient Alert Card.
- If your doctor feels that you are at risk for TB, you may be treated with medicines for TB before you are given Flixabi.

Tell your doctor straight away if you get signs of TB during treatment with Flixabi. Signs include persistent cough, weight loss, feeling tired, fever, night sweats.

### Hepatitis B virus (HBV)

- Tell your doctor if you are a carrier or if you have or have had hepatitis B before you are given Flixabi.
- Tell your doctor if you think you might be at risk of contracting HBV.
- Your doctor should test you for HBV.
- Treatment with TNF blockers such as Flixabi may result in reactivation of hepatitis B virus in patients who carry this virus, which can be life-threatening in some cases.

### Heart problems

- Tell your doctor if you have any heart problems, such as mild heart failure.
- Your doctor will want to closely monitor your heart function.

Tell your doctor straight away if you get new or worsening signs of heart failure during treatment with Flixabi. Signs include shortness of breath or swelling of your feet.

### Cancer and lymphoma

- Tell your doctor if you have or have ever had lymphoma (a type of blood cancer) or any other cancer before you are given Flixabi.
- Patients with severe rheumatoid arthritis, who have had the disease for a long time, may be at higher than average risk of developing lymphoma.
- Children and adults taking Flixabi may have an increased risk of developing lymphoma or another cancer.
- Some patients who have received TNF-blockers, including Flixabi have developed a rare type of cancer called Hepatosplenic T-cell Lymphoma. Of these patients, most were adolescent or young adult males and most had either Crohn's disease or ulcerative colitis. This type of cancer has usually resulted in death. Almost all patients had also received medicines known as azathioprine or 6-mercaptopurine in addition to TNF-blockers.
- Some patients treated with infliximab have developed certain kinds of skin cancer. If any changes in the appearance of the skin or growths on the skin occur during or after therapy, tell your doctor.
- Some women being treated for rheumatoid arthritis with infliximab have developed cervical cancer. For women taking infliximab including those over 60 years of age, your doctor may recommend that you continue to be regularly screened for cervical cancer.

#### Lung disease or heavy smoking

- Tell your doctor if you have a lung disease called Chronic Obstructive Pulmonary Disease (COPD) or if you are a heavy smoker, before you are given Flixabi.
- Patients with COPD and patients who are heavy smokers may have a higher risk of developing cancer with Flixabi treatment.

#### Nervous system disease

- Tell your doctor if you have or have ever had a problem that affects your nervous system before you are given Flixabi. This includes multiple sclerosis, Guillain-Barre syndrome, if you have fits or have been diagnosed with 'optic neuritis'.

Tell your doctor straight away if you get symptoms of a nerve disease during treatment with Flixabi. Signs include changes in your vision, weakness in your arms or legs, numbness or tingling in any part of your body.

#### Abnormal skin openings

- Tell your doctor if you have any abnormal skin openings (fistulae) before you are given Flixabi.

#### Vaccinations

- Talk to your doctor if you recently have had or are due to have a vaccine.
- You should not receive certain vaccines while using Flixabi.
- Certain vaccinations may cause infections. If you received Flixabi while you were pregnant, your baby may be at higher risk for getting such an infection up to six months after birth. It is important that you tell your baby's doctors and other health care professionals about your Flixabi use so they can decide when your baby should receive any vaccine, including live vaccines such as BCG (used to prevent tuberculosis). For more information see section on Pregnancy and breast-feeding.

#### Therapeutic infectious agents

- Talk to your doctor if you have recently received or are scheduled to receive treatment with a therapeutic infectious agent (such as BCG instillation used for the treatment of cancer).

#### Operations or dental procedures

- Tell your doctor if you are going to have any operations or dental procedures.
- Tell your surgeon or dentist performing the procedure that you are having treatment with Flixabi by showing them your Patient Alert Card.

#### **Children and adolescents**

The information above also applies to children and adolescents. In addition:

- Some children and teenage patients who have received TNF-blockers such as Flixabi have developed cancers, including unusual types, which sometimes resulted in death.
- More children taking Flixabi developed infections as compared to adults.
- Children should receive recommended vaccinations before starting Flixabi treatment.

If you are not sure if any of the above applies to you, talk to your doctor before you are given Flixabi.

#### **Other medicines and Flixabi**

Patients who have inflammatory diseases already take medicines to treat their problem. These medicines may cause side effects. Your doctor will advise you what other medicines you must keep using while you are having Flixabi.

Tell your doctor if you are using or have recently used any other medicines, including any other medicines to treat Crohn's disease, ulcerative colitis, rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis or psoriasis or medicines obtained without a prescription, such as vitamins and herbal medicines.

In particular, tell your doctor if you are using any of the following medicines:

- Medicines that affect your immune system.
- Kineret (anakinra). Flixabi and Kineret should not be used together.
- Orencia (abatacept). Flixabi and Orencia should not be used together.

While using Flixabi you should not receive live vaccines. If you were using Flixabi during pregnancy, tell your baby’s doctor and other health care professionals caring for your baby about your Flixabi use before the baby receives any vaccines.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before using Flixabi.

**Pregnancy and breast-feeding, and fertility**

- If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Flixabi should only be used during pregnancy if your doctor feels it is necessary for you.
- You should avoid getting pregnant when you are being treated with Flixabi and for 6 months after you stop being treated with it. Discuss the use of contraception during this time with your doctor.
- Do not breast-feed when you are being treated with Flixabi or for 6 months after your last treatment with Flixabi.
- If you received Flixabi during your pregnancy, your baby may have a higher risk for getting an infection.
- It is important that you tell your baby’s doctors and other healthcare professionals about your Flixabi use before your baby receives any vaccine. If you received Flixabi while pregnant, administration of BCG vaccine (used to prevent tuberculosis) to your baby within 6 months after birth may result in infection with serious complications, including death. Live vaccines such as BCG should not be given to your baby within 6 months after birth. For more information see section on vaccination.
- Severely decreased numbers of white blood cells have been reported in infants born to women treated with Flixabi during pregnancy. If your baby has continual fevers or infections, contact your baby’s doctor immediately.

**Driving and using machines**

Flixabi is not likely to affect your ability to drive or use tools or machines. If you feel tired or unwell after having Flixabi, do not drive or use any tools or machines.

**Flixabi contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially ‘sodium-free’.

**3. How Flixabi will be given**

**How Flixabi is given**

- Flixabi will be given to you by your doctor or nurse.
- Your doctor or nurse will prepare the Flixabi solution for injection.
- The Flixabi solution will be slowly injected (over a 2 hour period) into one of your veins. This will usually be in your arm. This is called an ‘intravenous infusion’ or drip. After the third treatment, your doctor may decide to give you Flixabi over a 1 hour period.
- You will be monitored while you are given Flixabi and also for 1 to 2 hours after.

**How much Flixabi is given**

- The doctor will decide your dose (in mg) and how often you will be given Flixabi. This will depend on your disease, weight and how well you respond to Flixabi.
- The table below shows how often you will usually have this medicine.

1 <sup>st</sup> treatment	0 weeks
2 <sup>nd</sup> treatment	2 weeks after your 1 <sup>st</sup> treatment
3 <sup>rd</sup> treatment	6 weeks after your 1 <sup>st</sup> treatment

Further treatments	Every 6 to 8 weeks depending on your disease
--------------------	--

### **Rheumatoid arthritis**

The recommended dose is 3 mg for every kg of body weight.

### **Psoriatic arthritis, ankylosing spondylitis (Bechterew's disease), psoriasis, ulcerative colitis and Crohn's disease**

The recommended dose is 5 mg for every kg of body weight.

### **Use in children and adolescents**

Flixabi should only be used in children and adolescents if they are being treated for Crohn's disease or ulcerative colitis. These children must be 6 years of age or older.

### **If you are given too much Flixabi**

As this medicine is being given by your doctor or nurse, it is unlikely that you will be given too much. There are no known side effects of having too much of Flixabi.

### **If you forget or miss your Flixabi infusion**

If you forget or miss an appointment to receive Flixabi, make another appointment as soon as possible.

If you have any further questions on the use of this medicine, ask your doctor.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. Most side effects are mild to moderate. However some patients may experience serious side effects and may require treatment. Side effects may also occur after your treatment with Flixabi has stopped.

### **Tell your doctor straight away if you notice any of the following:**

- **Signs of an allergic reaction** such as swelling of your face, lips, mouth or throat which may cause difficulty in swallowing or breathing, skin rash, hives, swelling of the hands, feet or ankles. An allergic reaction could happen within 2 hours of your injection or later. More signs of an allergic reaction that may happen up to 12 days after your injection include pain in the muscles, fever, joint or jaw pain, sore throat or headache
- **Signs of a heart problem** such as chest discomfort or pain, arm pain, stomach pain, shortness of breath, anxiety, lightheadedness, dizziness, fainting, sweating, nausea, vomiting, fluttering or pounding in your chest, a fast or a slow heartbeat, and/or swelling of your feet.
- **Signs of infection (including TB)** such as fever, feeling tired, (persistent) cough, shortness of breath, flu-like symptoms, weight loss, night sweats, diarrhoea, wounds, dental problems or burning when urinating
- **Signs of a lung problem** such as coughing, breathing difficulties or tightness in the chest
- **Signs of a nervous system problem (including eye problems)** such as fits, tingling or numbness in any part of your body, weakness in arms or legs, changes in eyesight such as double vision or other eye problems
- **Signs of a liver problem** such as yellowing of the skin or eyes, dark-brown coloured urine or pain in the upper right side of the stomach area, fever
- **Signs of an immune system disorder called lupus** such as joint pain or a rash on cheeks or arms that is sensitive to the sun
- **Signs of a low blood count** such as persistent fever, bleeding or bruising more easily or looking pale.

Tell your doctor straight away if you notice any of the above.

The known side effects of Flixabi include the following in groups of decreasing frequency:

**Very common side effects (affects more than 1 patient in 10)**

- Stomach pain, feeling sick
- Viral infections such as herpes or flu
- Upper respiratory infections such as sinusitis
- Headache
- Side effect due to an infusion
- Pain.

**Common side effects (affects 1 to 10 patients in 100)**

- Changes in how your liver works, increase in liver enzymes (shown in blood tests)
- Lung or chest infections such as bronchitis or pneumonia
- Difficult or painful breathing, chest pain
- Bleeding in the stomach or intestines, diarrhoea, indigestion, heartburn, constipation
- Nettle-type rash (hives), itchy rash or dry skin
- Balance problems or feeling dizzy
- Fever, increased sweating
- Circulation problems such as low or high blood pressure
- Bruising, hot flush or nosebleed, warm, red skin (flushing)
- Feeling tired or weak
- Bacterial infections such as blood poisoning, abscess or infection of the skin (cellulitis)
- Blood problems such as anaemia or low white blood cell count
- Swollen lymph nodes
- Depression, problems sleeping
- Eye problems, including red eyes and infections
- Fast heart beat (tachycardia) or palpitations
- Pain in the joints, muscles or back
- Urinary tract infection
- Psoriasis, skin problems such as eczema and hair loss
- Reactions at the injection site such as pain, swelling, redness or itching
- Chills, a build up of fluid under the skin causing swelling
- Feeling numb or having a tingling feeling.

**Uncommon side effects (affects 1 to 10 patients in 1000)**

- Shortage of blood supply, swelling of a vein
- Skin problems such as blistering, warts, abnormal skin colouration or pigmentation, or swollen lips
- Severe allergic reactions (e.g. anaphylaxis), an immune system disorder called lupus, allergic reactions to foreign proteins
- Wounds taking longer to heal
- Swelling of the liver (hepatitis) or gall bladder, liver damage
- Feeling forgetful, irritable, confused, nervous
- Eye problems including blurred or reduced vision, puffy eyes or sties
- New or worsening heart failure, slow heart rate
- Fainting
- Convulsions, nerve problems
- A hole in the bowel or blockage of the intestine, stomach pain or cramps
- Swelling of your pancreas (pancreatitis)
- Fungal infections such as yeast infection
- Lung problems (such as oedema)
- Fluid around the lungs (pleural effusion)
- Kidney infections
- Low platelet count, too many white blood cells
- Infections of the vagina.

**Rare side effects (affects 1 to 10 patients in 10,000)**

- A type of blood cancer (lymphoma)

- Your blood not supplying enough oxygen to your body, circulation problems such as narrowing of a blood vessel
- Inflammation of the lining of the brain (meningitis)
- Infections due to a weakened immune system
- Hepatitis B infection when you have had hepatitis B in the past
- Abnormal tissue swelling or growth
- Swelling of small blood vessels (vasculitis)
- Immune disorders that could affect the lungs, skin and lymph nodes (such as sarcoidosis)
- Lack of interest or emotion
- Serious skin problems such as toxic epidermal necrolysis, Stevens-Johnson Syndrome or erythema multiforme, skin problems such as boils
- Serious nervous system disorders such as transverse myelitis, multiple sclerosis-like disease, optic neuritis and Guillain-Barré syndrome
- Fluid in the lining of the heart (pericardial effusion)
- Serious lung problems (such as interstitial lung disease)
- Melanoma (a type of skin cancer).
- Cervical cancer
- Low blood counts, including a severely decreased number of white blood cells.

#### **Other side effects (the frequency is unknown)**

- Cancer in children and adults
- A rare blood cancer affecting mostly young people (hepatosplenic T-cell lymphoma)
- Liver failure
- Merkel cell carcinoma (a type of skin cancer).
- Worsening of a condition called dermatomyositis (seen as a skin rash accompanying muscle weakness).
- Temporary loss of sight during or within 2 hours of infusion
- The use of a 'live' vaccine may result in an infection caused by the 'live' viruses or bacteria contained in the vaccine (when you have a weakened immune system).

#### **Additional side effects in children and adolescents**

Children and adolescents who took infliximab for Crohn's disease showed some differences in side effects compared with adults who took infliximab for Crohn's disease. The side effects that happened more in children were: low red blood cells (anaemia), blood in stool, low white blood cells (leucopenia), redness or blushing (flushing), viral infections, low neutrophils which are white blood cells that fight infection (neutropenia), bone fracture, bacterial infection and allergic reactions of the breathing tract.

#### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Flixabi**

Flixabi will be stored by the health professionals at the hospital or clinic. The storage details should you need them are as follows:

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the label and the carton after "EXP". The expiry date refers to the last day of that month.
- Store in a refrigerator (2°C to 8°C).
- It is recommended that when Flixabi is prepared for infusion, it is used as soon as possible (within 3 hours). However, if the solution is prepared in germ-free conditions, it can be stored in a refrigerator at 2°C to 8°C for 24 hours.

- Do not use this medicine if it is discoloured or if there are particles present.

## 6. Contents of the pack and other information

### What Flixabi contains

- The active substance is infliximab. Each vial contains 100 mg of infliximab. After preparation each mL contains 10 mg of infliximab.
- The other ingredients are sucrose, polysorbate 80, monobasic sodium phosphate monohydrate and dibasic sodium phosphate heptahydrate.

### What Flixabi looks like and contents of the pack

Flixabi is supplied as a glass vial containing a powder for concentrate for solution for infusion. The powder is white.

Flixabi is produced in packs of 1 vial and in multipacks comprising 2, 3, 4, or 5 cartons, each containing 1 vial. Not all pack sizes may be marketed.

### Marketing Authorisation Holder

Samsung Bioepis UK Limited  
5th Floor, Profile West, 950 Great West Road  
Brentford  
Middlesex TW8 9ES  
United Kingdom

### Manufacturer

Biogen (Denmark) Manufacturing ApS  
Biogen Allé 1  
3400 Hillerød  
Denmark

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### België/Belgique/Belgien

Biogen Belgium NV/S.A  
Tél/Tel: + 32 (0)2 808 5947

#### Lietuva

Ewopharma AG Atstovybė  
Tel: +370 52 14 02 60

#### България

Ewopharma AG Representative Office  
Тел.: + 359 249 176 81

#### Luxembourg/Luxemburg

Biogen Belgium NV/SA  
Tél/Tel: +35 227 772 038

#### Česká republika

Biogen (Czech Republic) s.r.o.  
Tel: + 420 228 884 152

#### Magyarország

Biogen Hungary Kft.  
Tel.: + 36 (0)6 1 848 04 64

#### Danmark

Biogen (Denmark) A/S  
Tlf: + 45 78 79 37 53

#### Malta

Biogen Idec Limited (UK)  
Tel: + 356 27 78 15 79

#### Deutschland

Biogen GmbH  
Tel: + 49 (0)30 223 864 72

#### Nederland

Biogen Netherlands B.V.  
Tel: + 31 (0)20 808 02 70

#### Eesti

Ewopharma AG Eesti filiaal  
Tel: + 372 6 68 30 56

#### Norge

Biogen Norway AS  
Tlf: + 47 21 93 95 87

**Ελλάδα**

Biogen Idec Limited (UK)  
Τηλ: + 30 211 176 8555

**España**

Biogen Spain, S.L.  
Tel: + 34 931 790 519

**France**

Biogen France SAS  
Tél: + 33 (0)1 776 968 14

**Hrvatska**

Ewopharma d.o.o  
Tel: + 385 (0)1 777 64 37

**Ireland**

Biogen Idec (Ireland) Ltd.  
Tel: +353 (0)1 513 33 33

**Ísland**

Biogen Idec Limited (UK)  
Sími: + 354 800 9836

**Italia**

Biogen Italia s.r.l.  
Tel: + 39 (0)6 899 701 50

**Κύπρος**

Biogen Idec Limited (UK)  
Τηλ: + 357 22 00 04 93

**Latvija**

Ewopharma AG pārstāvniecība  
Tel: + 371 66 16 40 32

**Österreich**

Biogen Austria GmbH  
Tel: + 43 (0)1 267 51 42

**Polska**

Biogen Poland Sp. z o.o.  
Tel.: + 48 22 116 86 94

**Portugal**

Biogen Portugal Sociedade Farmacêutica,  
Unipessoal, Lda  
Tel: + 351 308 800 792

**România**

Ewopharma AG Representative Office  
Tel: + 40 377 881 045

**Slovenija**

Biogen Pharma d.o.o.  
Tel: + 386 (0)1 888 81 07

**Slovenská republika**

Biogen Slovakia s.r.o.  
Tel: + 421 (0)2 333 257 10

**Suomi/Finland**

Biogen Finland Oy  
Puh/Tel: + 358 (0)9 427 041 08

**Sverige**

Biogen Sweden AB  
Tel: +46 (0)8 525 038 36

**United Kingdom**

Biogen Idec Limited  
Tel: +44 (0)20 360 886 22

**This leaflet was last revised in {MM/YYYY}.**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.

This leaflet is available in all EU/EEA languages on the European Medicines Agency website.

-----  
**THE FOLLOWING INFORMATION IS INTENDED FOR HEALTHCARE PROFESSIONALS ONLY:**

***Instructions for use and handling – reconstitution, dilution and administration***

1. Calculate the dose and the number of Flixabi vials needed. Each Flixabi vial contains 100 mg infliximab. Calculate the total volume of reconstituted Flixabi solution required.
2. Under aseptic conditions, reconstitute each Flixabi vial with 10 ml of water for injections, using a syringe equipped with a 21-gauge (0.8 mm) or smaller needle. Remove flip-top from the vial and wipe the top with a 70% alcohol swab. Insert the syringe needle into the vial through the centre of the rubber stopper and direct the stream of water for injections to the glass wall of the vial. Gently swirl the solution by rotating the vial to dissolve the lyophilised powder. Avoid prolonged or vigorous agitation. **DO NOT SHAKE.** Foaming of the solution on reconstitution is not unusual. Allow the reconstituted solution to stand for 5 minutes. Check that the solution is colourless to light yellow and opalescent. The solution may develop a few fine translucent particles, as infliximab is a protein. Do not use if opaque particles, discolouration, or other foreign particles are present.
3. Dilute the total volume of the reconstituted Flixabi solution dose to 250 ml with sodium chloride 9 mg/ml (0.9%) solution for infusion. Do not dilute the reconstituted Flixabi solution with any other diluent. The dilution can be accomplished by withdrawing a volume of the sodium chloride 9 mg/ml (0.9%) solution for infusion from the 250-ml glass bottle or infusion bag equal to the volume of reconstituted Flixabi. Slowly add the total volume of reconstituted Flixabi solution to the 250-ml infusion bottle or bag. Gently mix.
4. Administer the infusion solution over a period of not less than the infusion time recommended. Use only an infusion set with an in-line, sterile, non-pyrogenic, low protein-binding filter (pore size 1.2 micrometre or less). Since no preservative is present, it is recommended that the administration of the solution for infusion is to be started as soon as possible and within 3 hours of reconstitution and dilution. When reconstitution and dilution are performed under aseptic conditions, Flixabi infusion solution can be used within 24 hours if stored at 2°C to 8°C. Do not store any unused portion of the infusion solution for reuse.
5. No physical biochemical compatibility studies have been conducted to evaluate the co-administration of Flixabi with other agents. Do not infuse Flixabi concomitantly in the same intravenous line with other agents.
6. Visually inspect Flixabi for particulate matter or discolouration prior to administration. Do not use if visibly opaque particles, discolouration or foreign particles are observed.
7. Any unused product or waste material should be disposed of in accordance with local requirements.